

RESEARCH ARTICLE

Open Access



LemurFaceID: a face recognition system to facilitate individual identification of lemurs

David Crouse^{1,7†}, Rachel L. Jacobs^{2*†}, Zach Richardson¹, Scott Klum^{1,8}, Anil Jain^{1*}, Andrea L. Baden^{3,4,5} and Stacey R. Tecot⁶

Abstract

Background: Long-term research of known individuals is critical for understanding the demographic and evolutionary processes that influence natural populations. Current methods for individual identification of many animals include capture and tagging techniques and/or researcher knowledge of natural variation in individual phenotypes. These methods can be costly, time-consuming, and may be impractical for larger-scale, population-level studies. Accordingly, for many animal lineages, long-term research projects are often limited to only a few taxa. Lemurs, a mammalian lineage endemic to Madagascar, are no exception. Long-term data needed to address evolutionary questions are lacking for many species. This is, at least in part, due to difficulties collecting consistent data on known individuals over long periods of time. Here, we present a new method for individual identification of lemurs (LemurFaceID). LemurFaceID is a computer-assisted facial recognition system that can be used to identify individual lemurs based on photographs.

Results: LemurFaceID was developed using patch-wise Multiscale Local Binary Pattern features and modified facial image normalization techniques to reduce the effects of facial hair and variation in ambient lighting on identification. We trained and tested our system using images from wild red-bellied lemurs (*Eulemur rubriventer*) collected in Ranomafana National Park, Madagascar. Across 100 trials, with different partitions of training and test sets, we demonstrate that the LemurFaceID can achieve $98.7\% \pm 1.81\%$ accuracy (using 2-query image fusion) in correctly identifying individual lemurs.

Conclusions: Our results suggest that human facial recognition techniques can be modified for identification of individual lemurs based on variation in facial patterns. LemurFaceID was able to identify individual lemurs based on photographs of wild individuals with a relatively high degree of accuracy. This technology would remove many limitations of traditional methods for individual identification. Once optimized, our system can facilitate long-term research of known individuals by providing a rapid, cost-effective, and accurate method for individual identification.

Keywords: Animal biometrics, Conservation, *Eulemur rubriventer*, Linear discriminant analysis, Mammal, Multiscale local binary pattern, Pelage, Photograph, Primate

Background

Most research on the behavior and ecology of wild animal populations requires that study subjects are individually recognizable. Individual identification is necessary to ensure unbiased data collection and to account for individual variation in the variables of interest. For short-term

studies, researchers may rely on unique methods for identification based on conspicuous natural variation among individuals at the time of data collection, such as differences in body size and shape or the presence of injuries and scars. These methods may or may not allow for identification of individuals at later dates in time. To address many evolutionary questions, however, it is necessary to collect data on known individuals over long periods of time [1]. Indeed, longitudinal studies are essential for characterizing life history parameters, trait heritability, and fitness effects (reviewed in [1]). Consequently, they are invaluable for identifying the demographic and evolutionary processes influencing wild animal populations [1].

* Correspondence: rachel_jacobs@gwu.edu; jain@cse.msu.edu

†Equal contributors

²Department of Anthropology, Center for the Advanced Study of Human Paleobiology, The George Washington University, Washington, DC, USA

¹Department of Computer Science and Engineering, Michigan State University, East Lansing, MI, USA

Full list of author information is available at the end of the article



Unfortunately, longitudinal monitoring can be challenging, particularly for long-lived species. One of the primary challenges researchers face is establishing methods for individual identification that allow multiple researchers to collect consistent and accurate demographic and behavioral data over long periods of time (in some cases several decades). Current methods for individual identification often involve either capturing and tagging animals with unique identifiers, such as combinations of colored collars and/or tags [2–5], or taking advantage of natural variation in populations (e.g., scars, skin and pelage patterns) and relying on researchers' knowledge of individual differences [6–9]. The former method (or a combination of the two methods) has been used in some of the best established long-term field studies, such as the St. Kilda Soay Sheep and Isle of Rum Red Deer Projects [2, 3], as well as the Wytham Tit and Galápagos Finch Projects [4, 5]. Because they have long-term (multi-generation) data on known individuals, these projects have contributed substantially to the field of evolutionary biology by documenting how and why populations change over time (e.g., [10–13]).

Similar methods involving capturing and collaring have been used in many longitudinal studies of wild primates, such as owl monkeys [14], titi monkeys [15], colobines [16], and in particular, many Malagasy lemurs [17–20]. Through the long-term monitoring of individuals, many of these studies have provided important data on longevity, lifetime reproductive success, and dispersal patterns [15, 17, 18, 20–23].

Despite its utility for many longitudinal studies, the tagging process might sometimes be inappropriate or otherwise impractical. Tagging often requires that study subjects be captured via mist netting or in nest boxes (for birds) [4, 5], trapping (e.g., Sherman traps or corrals for some mammals) [2, 3, 24], and, in the case of some larger mammals, including many primates, darting via blow gun or air rifle [10, 25–27]. Capturing has several advantages, such as enabling data to be collected that would otherwise be impossible (e.g., blood samples, ectoparasites), but it can also be expensive, often making it unfeasible for studies with large sample sizes and/or those conducted over large spatial and temporal scales. Furthermore, capturing and tagging may pose additional risks to already threatened species. For example, such methods have been shown in some cases to cause acute physiological stress responses [16], tissue damage [28] and injury (e.g., broken bones, paralysis) [29], as well as disrupt group dynamics, and pose risks to reproduction, health, and even life [29–32].

An alternative method for individual identification relies on researcher knowledge of variation in individual appearances. It is less invasive and removes some of the potential risks associated with capturing and tagging.

Such methods have been successfully used in long-term studies of elephants, great apes, and baboons (among others) and have provided similarly rich long-term datasets that have been used to address demographic and evolutionary questions [6–9]. However, this method is more vulnerable to intra- and inter-observer error and thus can require substantial training. Moreover, for research sites involving multiple short-term studies in which researchers may use different methods for individual identification, it can be difficult to integrate data [33]. Additionally, long-term research is often hindered by disruptions to data collection (e.g., between studies, due to lack of research funds, political instability [1]). These breaks can result in lapses of time during which no one is present to document potential changes to group compositions and individual appearances, which can also complicate integrating data collected at different time points.

Under such circumstances, projects would benefit from a database of individual identifications, as well as a rapid method for identifying individuals that requires little training and can be used across different field seasons and researchers. The field of animal biometrics offers some solutions [34]. For example, some methods that have shown promise in mammalian (among other) research, including studies of cryptic animals, combine photography with computer-assisted individual identification programs to facilitate long-term systematic data collection (e.g., cheetahs: [35]; tigers: [36]; giraffes: [37]; zebras: [38]). These methods use quantifiable aspects of appearances to identify individuals based on probable matches in the system [34]. Because assignments are based on objective measures, these methods can minimize intra- and inter-observer error and facilitate integrating data collected across different studies [34]. At the same time, in study populations with large sample sizes, researchers might be limited in the number of individuals known on-hand. Computer-assisted programs can facilitate processing data to rapidly identify individuals when datasets are large, which reduces the limitations on sample size/scale imposed by the previous methods [34].

Despite their potential utility, such methods have not been incorporated in most studies of wild primates, and, particularly in the case of wild lemur populations, even with several drawbacks, capture and collar methods remain common [17–20]. As a result, multi-generation studies of lemur populations that incorporate individual identification are limited.

Here we present a method in development for non-invasive individual identification of wild lemurs that can help mitigate some of the disadvantages associated with other methods, while also facilitating long-term research (Table 1). Our system, called LemurFaceID, utilizes

Table 1 Individual identification methods

Method	Advantages	Disadvantages
Tagging/Collaring	Systematic across studies; opportunities to collect data that require animal to be in hand; precise location of animal known at all times (using GPS collar)	Invasive; poses risks to animals; expensive; less feasible for studies requiring large sample sizes; individual IDs may be unknown with loss of tag/collar
Manual identification based on physical variation	Non-invasive, low cost	Substantial training required; IDs may differ across studies/researchers; prone to intra- and inter-observer error; time-consuming for large sample sizes when individuals are not recognized instantly (e.g., manual comparisons of photographs are required)
Face recognition	Systematic across studies; non-invasive; minimal user training; reduces time to make identifications when datasets are large allowing for increased sample size/scale	Requires large dataset for development; currently requires partial knowledge of individual IDs; individual IDs may be unknown to the researcher if the system is unavailable for use

computer facial recognition methods, developed by the authors specifically for lemur faces, to identify individual lemurs based on photographs collected in wild populations [39].

Facial recognition technology has made great strides in its ability to successfully identify humans [40], but this aspect of computer vision has much untapped potential. Facial recognition technology has only recently expanded beyond human applications. While there has been limited work with non-human primates [41, 42], to our knowledge, facial recognition technology has not been applied to any of the >100 lemur species. However, many lemurs possess unique facial features, such as hair/pelage patterns, that make them appropriate candidates for applying modified techniques developed for human facial recognition (Fig. 1).

We focus this study on the red-bellied lemur (*Eulemur rubriventer*). Males and females in this species are sexually dichromatic with sex-specific variation in facial patterns ([43]; Fig. 2). Males exhibit patches of white skin around the eyes that are reduced or absent in females. In addition, females have a white ventral coat (reddish-brown in males) that variably extends to the neck and face. Facial patterns are individually variable, and the authors have used this variation to identify individuals in wild populations, but substantial training was required. Since the 1980s, a population of red-bellied lemurs has been studied in Ranomafana National Park, Madagascar [44–47], but because researchers used different methods for individual identification, gaps between studies make it difficult to integrate data. Consequently, detailed data on many life history parameters for this





Fig. 2 Red-bellied lemurs. The individual on the right is female, and the individual on the left is male

species are lacking. A reliable individual identification method would help provide these critical data for understanding population dynamics and addressing evolutionary questions.

In this paper we report the method and accuracy results of LemurFaceID, as well as its limitations. This system uses a relatively large photographic dataset of known individuals, patch-wise Multiscale Local Binary Pattern (MLBP) features, and an adapted Tan and Triggs [48] approach to facial image normalization to suit lemur face images and improve recognition accuracy.

Our initial effort (using a smaller dataset) was focused on making parametric adaptations to a face recognition system designed for human faces [49]. This system used both MLBP features and Scale Invariant Feature Transform (SIFT) features [50, 51] to characterize face images. Our initial effort exhibited low performance in recognition of lemur faces (73% rank-1 recognition accuracy). In other words, for a given query, the system reported the highest similarity between the query and the true match in the database only 73% of the time. Examination of the system revealed that the SIFT features were sensitive to local hair patterns. As matting of hair changed from image to image, the features changed substantially and therefore reduced match performance. The high dimensionality of the SIFT features also may have led to overfitting and slowing of the recognition process. Because of this, the use of SIFT features was abandoned in the final recognition system.

While still adapting methods originally developed for humans, LemurFaceID is specifically designed to handle lemur faces. We demonstrate that the LemurFaceID system identifies individual lemurs with a level of accuracy that suggests facial recognition technology is a potential useful tool for long-term research on wild lemur populations.

Methods

Data collection

Study species

Red-bellied lemurs (*Eulemur rubriventer*) are small to medium-sized (~2 kg), arboreal, frugivorous primates, and they are endemic to Madagascar's eastern rainforests [46, 52] (Fig. 3a). Despite their seemingly widespread distribution, the rainforests of eastern Madagascar have become highly fragmented [53], resulting in an apparent patchy distribution for this species. It is currently listed by the IUCN as Vulnerable with a decreasing population trend [54].


Study site

Data collection for this study was concentrated on the population of red-bellied lemurs in Ranomafana National Park (RNP). RNP is approximately 330 km² of montane rainforest in southeastern Madagascar [22, 55] (Fig. 3b). Red-bellied lemurs in RNP have been the subjects of multiple research projects beginning in the 1980s [44–47].

Dataset

Our dataset consists of 462 images of 80 red-bellied lemur individuals. Each individual had a name (e.g., Avery) or code (e.g., M9VAL) assigned by researchers when it was first encountered. Photographs of four individuals are from the Duke Lemur Center in North Carolina, while the remainder are from individuals in RNP in Madagascar. The number of images (1–21) per individual varies. The dataset only includes images that contain a frontal view of the lemur's face with little to no obstruction or occlusion. The dataset comprises images with a large range of variation; these include images with mostly subtle differences in illumination and focus (generally including subtle differences in gaze; ~25%), as well as images with greater variation (e.g., facial orientation, the presence of small obstructions, illumination and shadows; ~75%). Fig. 4 contains a histogram of the number of images available per individual. Amateur photographers captured photos from RNP using a Canon EOS Rebel T3i with 18–55 and 75–300 mm lenses. Lemurs were often at heights between 15–30 m, and photos were taken while standing on the ground. Images from the Duke Lemur Center were captured with a Google Nexus 5 or an Olympus E-450 with a 14–42 mm lens. Lemurs were in low trees (0–3 m), on the ground, or in enclosures, and photos were taken while standing on the ground.

The majority of images taken in Madagascar were captured from September 2014 to March 2015, though some individuals had images captured as early as July 2011. Images from the Duke Lemur Center were captured in July 2014. Due to the longer duration of the image collection in Madagascar, there was some difficulty establishing whether certain individuals encountered in



2014 had been encountered previously. In three cases, there are photographs in the dataset labeled as belonging to two separate individuals that might be of the same individual. These images were treated as belonging to separate individuals when partitioning the dataset for experiments, but if images that might belong to a single individual were matched together, it was counted as a successful match. Figure 5 illustrates the facial similarities and variations present in the dataset. Figure 5a illustrates the similarities and differences between the 80 wild individuals (inter-class similarity), while Fig. 5b shows different images of the same individual (intra-class variability). In addition to the database of red-bellied lemur individuals, a database containing lemurs of other species was assembled. This

database includes 52 images of 31 individuals from Duke Lemur Center and 138 images of lemurs downloaded using an online image search through Google Images. We used only those images with no apparent copyrights. These images were used to expand the size of the gallery for lemur identification experiments.

Recognition system

Figure 6 illustrates the operation of our recognition system (LemurFaceID). This system was implemented using the OpenBR framework (openbiometrics.org; [56]).

Image pre-processing

Eye locations have been found to be critical in human face recognition [40]. The locations of eyes are critical to normalizing the facial image for in-plane rotation. We were unable to design and train a robust eye detector for lemurs because our dataset was not sufficiently large to do so. For this reason, we used manual eye location. Prior to matching, the user marks the locations of the lemur’s eyes in the image. Using these two points, with the right eye as the center, a rotation matrix M is calculated to apply an affine transformation to align the eyes horizontally. Let lex , ley , rex , and rey represent the x and y coordinates of the left and right eyes, respectively. The affine matrix is defined as:






Fig. 5 Variation in lemur face images. **a** Inter-class variation. **b** Intra-class variation. Some images in this figure are modified (i.e., cropped) versions of images that have been previously published in [74] under a CC BY License

$$M = \begin{bmatrix} 0 & 0 & rex \\ 0 & 0 & rey \\ 0 & 0 & 1 \end{bmatrix} \times \begin{bmatrix} \cos(\theta) & -\sin(\theta) & 0 \\ \sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 1 \end{bmatrix} \times \begin{bmatrix} 0 & 0 & -rex \\ 0 & 0 & -rey \\ 0 & 0 & 1 \end{bmatrix}$$

$$\theta = \text{atan}\left(\frac{ley-rey}{lex-rer}\right)$$

The input image is rotated by the matrix M and then cropped based on the eye locations. Rotation is applied

prior to cropping so that the area cropped will be as accurate as possible. The Inter-Pupil Distance (IPD) is taken as the Euclidean distance between the eye points. The image is cropped so that the eyes are $\frac{IPD}{2}$ pixels from the nearest edge and $0.7 \times IPD$ pixels from the top edge, with a total dimension of $IPD \times 2$ pixels square. This image is then resized to the final size of 104×104 pixels, which facilitates the patch-wise feature extraction




Fig. 6 Flowchart of LemurFaceID. Linear discriminant analysis (LDA) is used for reducing feature vector dimensionality to avoid overfitting

scheme described below. This process is illustrated in Fig. 7. Following rotation and cropping, the image is converted to gray-scale and normalized. Although individual lemurs do show variation in pelage/skin coloration, we disregard color information from the images. In human face recognition studies, skin color is known to be sensitive to illumination conditions and therefore is not considered to be a reliable attribute [57, 58].

Since the primary application of the LemurFaceID system is to identify lemurs from photos taken in the wild, the results must be robust with respect to illumination variations. To reduce the effects of ambient illumination on the matching results, a modified form of the illumination normalization method outlined by Tan and Triggs [48] is applied. The image is first convolved with a Gaussian filter with $\sigma = 1.1$, and is then gamma corrected ($\gamma = 0.2$). A Difference of Gaussians (DoG) operation [48] (with parameters σ_1 and σ_2 corresponding to the standard deviations of the two Gaussians) is subsequently performed on the image. This operation

eliminates small-scale texture variations and is traditionally performed with $\sigma_1 = 1$ and $\sigma_2 = 2$. In the case of lemurs, there is an ample amount of hair with a fine texture that varies from image to image within individuals. This fine texture could confuse the face matcher, as changes in hair orientation would result in increased differences between face representations. To reduce this effect in the normalized images, σ_1 is set to 2. The optimal value of σ_2 was empirically determined to be 5. The result of this operation is then contrast equalized using the method outlined in Tan and Triggs [48], producing a face image suitable for feature extraction. Figure 8 illustrates a single lemur image after each pre-processing step.

Feature extraction

Local Binary Pattern (LBP) representation is a method of characterizing local textures in a patch-wise manner [50]. Each pixel in the image is assigned a value based on its relationship to the surrounding pixels, specifically based on whether each surrounding pixel is darker than





Fig. 7 Eye selection, rotation, and cropping of a lemur image



the central pixel or not. Out of the 256 possible binary patterns in a 3×3 pixel neighborhood, 58 are defined as uniform (having no more than 2 transitions between “darker” and “not darker”) [50]. The image is divided into multiple patches (which may or may not overlap), and for each patch a histogram of the patterns is developed. Each of the 58 uniform patterns occupies its own bin, while the non-uniform patterns occupy a 59th bin [50]. This histogram makes up a 59-dimensional feature vector for each patch. In our recognition system, we use 10×10 pixel patches, overlapping by 2 pixels on a side. This results in 144 total patches for the 104×104 face image.


Multi-scale Local Binary Pattern (MLBP) features are a variation on LBP which use surrounding pixels at different radii from the central pixel [50], as shown in Fig. 9.


For this application, we used radii of 2, 4, and 8 pixels. Therefore, each patch generates 3 histograms, one per radius, each of which is normalized, and then concatenated and normalized again, both times by L2 norm. This process results in a 177-dimensional feature vector for each 10×10 patch. Figure 10 shows an example of three face images of the same individual with an enlarged grid overlaid. As demonstrated by the highlighted areas, patches from the same area in each image will be compared in matching.

To extract the final feature vector, linear discriminant analysis (LDA) is performed on the 177-dimensional feature vector for each patch. LDA transforms the feature vector into a new, lower-dimensional feature vector such that the new vector still captures 95% of the variation between individuals, while minimizing the amount of variation between images of the same individual. For this transformation to be robust, a large training set of lemur face images is desirable. LDA is trained on a per-patch basis to limit the size of the feature vectors considered. The resulting vectors for all the patches are then concatenated and normalized to produce the final feature vector for the image. Because each patch undergoes its own dimensionality reduction, the final dimensionality of the feature vector will vary from one training set to another. The LemurFaceID system reduces the mean size of the resultant image features from 396,850 dimensions to 7,305 dimensions.

Face matching

In preparation for matching two lemur faces, a gallery (a database of face images and their identities against which a query is searched) is assembled containing feature representations of multiple individual lemurs. The Euclidean distance d between feature vectors of a query image and each image in the gallery is calculated. The final similarity metric is defined as $[1 - \log(d + 1)]$; higher values indicate more similar faces. A query can consist of 1 or more images, all of which must be of the same lemur. For each query image, the highest similarity score for each individual represents that individual’s match score. The mean of these scores, over multiple query





images, is calculated to obtain the final individual scores. The top five ranking results (i.e., individuals with the 5 highest scores) are presented in descending order. We evaluated LemurFaceID systems' recognition performance with queries consisting of 1 and 2 images.

Figure 11a shows match score histograms for genuine (comparing 2 instances of the same lemur) vs. impostor (comparing 2 instances of different lemurs) match scores with 1 query image. Figure 11b shows score histograms with fusion of 2 query images. Note that the overlap between genuine and impostor match score histograms is substantially reduced by the addition of a second query image.

Statistical analysis

We evaluated the accuracy of the LemurFaceID system by conducting 100 trials over random splits of the lemur face dataset (462 images of 80 red-bellied lemurs) that we collected. To determine the response of the recognition system to novel individuals, the LDA dimensionality reduction method must be trained on a different set of individuals (i.e., training set) from those used to evaluate matching performance (known as the test set). To satisfy this condition, the dataset was divided into training and testing sets via random split. Two-thirds of the 80 individuals (53 individuals) were designated as the training set, while the remainder (27 individuals) comprised the test set. In the test set, two-thirds of the images for each individual were assigned to the system database (called the 'gallery' in human face recognition literature) and the remaining images were assigned as queries (called the 'probe' in human face recognition literature). Individuals with fewer than 3 images were placed only in the gallery. The gallery was then expanded to include a secondary dataset of other species to increase its size.

Testing was performed in open-set and closed-set identification scenarios. Open-set mode allows for conditions encountered in the wild, where lemurs (query images) may be encountered that have not been seen before (i.e., individuals are not present in the system

database). Queries whose fused match score is lower than a certain threshold are classified as containing a novel individual. Closed-set mode assumes that the query lemur (lemur in need of identification) is represented in the gallery and may be useful for identifying a




Fig. 11 Histograms of genuine (correct match) vs. impostor (incorrect match) scores. **a** Results with only one query image (4,265 genuine, 831,583 impostor). **b** Results with 2 query images (4,317 genuine, 841,743 impostor)

lemur in situations where the system is guaranteed to know the individual, such as in a captive colony.

For open-set testing, one-third of the red-bellied lemur individuals in the gallery were removed. Their corresponding images in the probe set therefore made up the set of novel individuals. For open-set, the mean gallery size was 266 images, while for closed-set the mean size was 316 images. Across all trials of the LemurFaceID system, the mean probe size was 42 images.

Results

Results of the open-set performance of LemurFaceID are presented in Fig. 12, which illustrates the Detection and Identification Rate (DIR) against the False Accept Rate (FAR). DIR is calculated as the proportion of non-novel individuals that were correctly identified at or below a given rank. FAR is calculated as the number of novel individuals incorrectly matched to a gallery individual at or below a given rank. In general, individuals are correctly identified >95% of the time at rank 5 or higher regardless of FAR, but DIR is lower (<95%) at rank 1, only approaching 95% when FAR is high (0.3).

Rank 1 face matching results for closed-set operation are reported in Table 2, and the Cumulative Match Characteristic (CMC) curves for 1-image query and 2-

image fusion (combining matching results for the individual query images) are shown in Fig. 13. This plot shows the proportion of correct identifications at or below a given rank. The mean percentage of correct matches (i.e., Mean True Accept Rate) increases when 2 query images are fused; individuals are correctly identified at Rank 1 $98.7\% \pm 1.81\%$ using 2-image fusion compared to a Rank 1 accuracy of $93.3\% \pm 3.23\%$ when matching results for a single query image are used.

Discussion

Our initial analyses of LemurFaceID suggest that facial recognition technology may be a useful tool for individual identification of lemurs. This method represents, to our knowledge, the first system for machine identification of lemurs by facial features. LemurFaceID exhibited a relatively high level of recognition accuracy (98.7%; 2-query image fusion) when used in closed-set mode (i.e., all individuals are present in the dataset), which could make this system particularly useful in captive settings, as well as wild populations with low levels of immigration from unknown groups. Given the success of LemurFaceID in recognizing individual lemurs, this method could also allow for a robust species recognition system, which would be useful for presence/absence studies.




Table 2 Face matcher evaluation results (Rank 1, closed-set)

Method	Mean (TAR)	SD
Baseline system	81.5%	6.68%
2 query images	98.7%	1.81%
1 query image	93.3%	3.23%

True Accept Rate (TAR) is the percentage of correct matches. Standard deviation (SD) is computed over 100 random splits. The LemurFaceID system is also compared to the earlier (i.e., "Baseline") system (using SIFT) for comparison


The accuracy of our system was lower using open-set mode (i.e., new individuals may be encountered) where, regardless of the False Accept Rate (FAR), non-novel individuals were correctly identified at rank 1 less than 95% of the time and less than 85% of the time given a FAR of 0. These numbers are expected to improve with a larger dataset of photographs and individuals. In our current sample, we also included photographs exhibiting only subtle variation between images. Given that the

ultimate goal of LemurFaceID is to provide an alternative, non-invasive identification method for long-term research, it will also be important to test its accuracy using a larger dataset that includes only photographs with large variation (e.g., collected across multiple, longer-term intervals).

We also note that our system focuses specifically on classifying individuals using a dataset of known individuals in a population. Such a tool can be particularly useful for maintaining long-term research on a study population. This approach differs, however, from another potential application of face recognition methods, which would be to identify the number of individuals from a large image dataset containing unknown individuals only (i.e., clustering) [59, 60]. The addition of a clustering technique could allow for more rapid population surveys or facilitate the establishment of new study sites, but such techniques can be challenging as clustering accuracy is expected to be lower than the classification accuracy [59, 60]. That said, in future work, the feature extraction and scoring system of LemurFaceID could potentially be combined with clustering techniques for segmenting datasets of unknown individuals.

Despite some current limitations, LemurFaceID provides the groundwork for incorporating this technology into long-term research of wild lemur populations, particularly of larger-bodied (>2 kg) species. Moving forward, we aim to 1) expand our photographic database, which is necessary to automate the lemur face detector and eye locator, 2) increase open-set performance by improving the feature representation to provide better separation between scores for in-gallery and novel individuals, and 3) field test the system to compare the classification accuracy of LemurFaceID with that of experienced and inexperienced field observers. Once optimized, a non-invasive, computer-assisted program for individual identification in lemurs has the potential to mitigate some of the challenges faced by long-term research using more traditional methods.

For example, facial recognition technology would remove the need to artificially tag individuals, which removes potential risks to animals associated with capturing and collaring; some of these risks, including injury, occur more frequently in arboreal primates [29]. At the same time, many costs incurred using these techniques are removed (e.g., veterinary services, anesthesia), as are potential restrictions on the number of individuals available for study (e.g., local government restrictions on captures). More traditional non-invasive techniques that rely on researchers' knowledge of natural variation can be similarly advantageous, but facial recognition programs can help ensure that data are collected consistently across multiple researchers. That said, we would



not recommend researchers become wholly reliant on computer programs for individual identification of study subjects, but training multiple researchers to accurately recognize hundreds of individuals is time-consuming and costly, as well as potentially unrealistic. Facial recognition technology can facilitate long-term monitoring of large populations by removing the need for extensive training, or potentially accelerate training by making phenotypic differences more tangible to researchers and assistants. Moreover, in studies with large sample sizes where immediate recognition of all individuals might be impossible, facial recognition technology can process data more quickly. For example, LemurFaceID takes less than one second to recognize a lemur (using a quad core i7 processor), which will save time identifying individuals when manual comparisons of photographs/descriptions are necessary.

Ultimately then, LemurFaceID can help expand research on lemur populations by providing a method to systematically identify a large number of individuals over extended periods of time. As is the case with other long-term studies of natural populations, this research has the potential to provide substantial contributions to evolutionary biology [1]. More specifically, lemurs are an endemic mammalian lineage that evolved in Madagascar beginning >50 million years ago [61]. Over time, they have greatly diversified with >100 species recognized today [43]. They occupy diverse niches (e.g., small-bodied, nocturnal gummivores; arrhythmic frugivores; large-bodied, diurnal folivores) across Madagascar's varied habitats (e.g., rainforests; spiny, dry forest) [43], and they have recently (in the last ~2,000 years) experienced extensive ecological change owing largely to human impact [62]. Accordingly, this mammalian system provides unique opportunities for studying ecological and evolutionary pressures impacting wild populations.

Data obtained from longitudinal studies of lemurs can also aid in conservation planning and management for this highly endangered group of mammals. Demographic structure and life history parameters documented from long-term research can provide insights into the causes of population change and be used to model extinction risk [63–65]. LemurFaceID also has potential for more direct applications to conservation. One notable threat to lemurs [66, 67], as well as many other animal species [68, 69], is live capture of individuals for the pet trade. LemurFaceID could provide law enforcement, tourists, and researchers with a tool to rapidly report sightings and identify captive lemurs (species and individuals). A database of captive lemurs can help with continued monitoring to determine if individuals remain constant over time.

Importantly, the face recognition methods we developed for LemurFaceID could be useful for individual identification in other primates, as well as other non-

primate species, especially those with similarly variable facial pelage/skin patterns (e.g., bears, red pandas, raccoons, sloths). Furthermore, as camera trapping has become increasingly useful for population monitoring of many cryptic species (e.g., [70, 71]), our facial recognition technology could be potentially incorporated into long-term, individual-based studies conducted remotely. That said, it will be necessary to make unique modifications to methods for different lineages.

To illustrate this point, recent publications also have explored the area of facial recognition for primates. For example, Loos and Ernst's [41] system for recognizing chimpanzees has a similar approach to pre-processing as LemurFaceID, but they use a different illumination normalization method and correct for greater difference in perspective. In feature extraction, their use of speeded-up robust features (SURF), a gradient-based feature similar to SIFT, underscores the difference in lemur and chimpanzee faces, namely the lack of hair/fur in chimpanzees to confound the directionality of the features [41]. Their selection of Gabor features also reflects the relative lack of hair, as such indicators of edgeness would exhibit significantly more noise in lemurs [72]. More recently, Freytag et al. [73] were able to improve upon recognition accuracy of chimpanzees by applying convolutional neural network (CNN) techniques. Their results identify CNNs to be a promising direction of animal face recognition research, but such methods also require datasets that are orders of magnitude larger than our current dataset [73]. Thus, although they are beyond the scope of this study, CNNs could be an interesting avenue for future research in lemur face recognition.

In contrast to these approaches, Allen and Higham [42] use a biologically-based model for identification of guenons. Their feature selection is based on guenon vision models, using the dimensions of facial spots to identify species and individuals [42]. While *E. rubriventer* individuals also possess prominent facial spots, these are not common across different lemur species and therefore unsuitable for use in our system. The wide variety of approaches used underscores that there is no "one size fits all" approach to animal facial recognition, but once developed, this technology has the potential to facilitate long-term research in a host of species, expand the types of research questions that can be addressed, and help create innovative conservation tools.

Conclusions

Our non-invasive, computer-assisted facial recognition program (LemurFaceID) was able to identify individual lemurs based on photographs of wild individuals with a relatively high degree of accuracy. This technology would remove many limitations of traditional methods for individual identification of lemurs. Once optimized,

our system can facilitate long-term research of known individuals by providing a rapid, cost-effective, and accurate method for individual identification.

Abbreviations

CMC: Cumulative match characteristic; CNN: Convolutional neural network; DIR: Detection and Identification Rate; FAR: False accept rate; IPD: Inter-pupil distance; LBP: Local binary pattern; LDA: Linear discriminant analysis; MLBP: Multiscale local binary pattern; RNP: Ranomafana National Park; SIFT: Scale invariant feature transform; SURF: Speeded-up robust features; TAR: True accept rate

Acknowledgements

Logistics and permissions for research in Madagascar were facilitated by Benjamin Andriamihaja and MICET, Ministre des Eaux et Forets, Madagascar National Parks, Eileen Larney and the Centre ValBio, and the University of Antananarivo. We would like to thank Samantha Ambler, Caroline Angyal, Alicia S. Arroyo, Bashira Chowdhury, Joseph Falinomenjanahary, Sheila Holmes, Jean Pierre Lahitsara, Avery Lane, Natalee Phelps, Aura Raulo, Soafaniry Razanajatovo, and Jean Baptiste Velontsara for their contribution to collecting face images of lemurs, as well as multiple students, volunteers, and research technicians for assisting with data collection. Finally, we thank the Duke Lemur Center (DLC) staff for logistical support during data collection at the DLC. This is DLC publication number 1336.

Funding

This research was supported by funds from the American Association of Physical Anthropologists to SRT, National Science Foundation (DDIG, BCS 1232535) to RLJ, The Leakey Foundation to RLJ, SRT, and ALB, The Wenner-Gren Foundation to RLJ, Rowe-Wright Primate Fund to RLJ, SRT, and ALB, Stony Brook University to RLJ, IDEAWILD to RLJ and SRT, University of Arizona to SRT, Hunter College-CUNY to ALB, and Michigan State University to AJ, DC, and SK. Funders had no role in the design of the study, data collection, analysis, and interpretation, or preparation of the manuscript.

Availability of data and materials

The lemur images and source code for LemurFaceID, including instructions for its use, are available online through Michigan State University's Biometrics Research Group (http://biometrics.cse.msu.edu/Publications/Databases/MSU_LemurFaceID/).

Authors' contributions

RLJ and SRT conceived of the project. RLJ, SRT, ALB, DC, and ZR acquired data for the project. DC, SK, and AJ conceived and designed the recognition system and experiments. DC, ZR, SK, and AJ performed the experiments and analyzed data. RLJ, SRT, AJ, ALB, and DC drafted the manuscript. All authors read and approved of the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Not applicable.

Ethics approval and consent to participate

Data were collected in accordance with and approved by institutional committees (Stony Brook University IACUC: 2010/1803, 2011/1895; University of Arizona IACUC: 13-470) and Madagascar National Parks.

Author details

¹Department of Computer Science and Engineering, Michigan State University, East Lansing, MI, USA. ²Department of Anthropology, Center for the Advanced Study of Human Paleobiology, The George Washington University, Washington, DC, USA. ³Department of Anthropology, Hunter College, City University of New York, New York, NY, USA. ⁴The Graduate Center of City University of New York, New York, NY, USA. ⁵The New York Consortium in Evolutionary Primatology (NYCEP), New York, NY, USA. ⁶School of Anthropology, The University of Arizona, Tucson, AZ, USA. ⁷Present address: Samsung Semiconductor, Inc., San Jose, CA, USA. ⁸Present address: Noblis, Inc., Annandale, VA, USA.

Received: 8 August 2016 Accepted: 29 December 2016

Published online: 17 February 2017

References

- Clutton-Brock T, Sheldon BC. Individuals and populations: the role of long-term, individual-based studies of animals in ecology and evolutionary biology. *Trends Ecol Evol.* 2010;25:562–73.
- Clutton-Brock T, Pemberton J. Soay sheep: dynamics and selection in an island population. Cambridge: Cambridge University Press; 2004.
- Clutton-Brock TH. Red deer: the behaviour and ecology of two sexes. Chicago: University of Chicago Press; 1982.
- Lack D, Gibb J, Owen DF. Survival in relation to brood-size in tits. *J Zool.* 1957;128:313–26.
- Grant PR, Grant BR. 40 years of evolution: Darwin's finches on Daphne major island. Princeton: Princeton University Press; 2014.
- Moss CJ. The demography of an African elephant (*Loxodonta africana*) population in Amboseli, Kenya. *J Zool.* 2001;255:145–56.
- Murray CM, Stanton MA, Wellens KR, Santymire RM, Heintz MR, Lonsdorf EV. Maternal effects on offspring stress physiology in wild chimpanzees. *Am J Primatol.* 2016, in press. DOI:10.1002/ajp.22525.
- Wich SA, Utami-Atmoko SS, Setia TM, Rijksen HD, Schürmann C, van Hooff JARAM, van Schaik CP. Life history of wild Sumatran orangutans (*Pongo abelii*). *J Hum Evol.* 2004;47:385–98.
- Alberts SC, Altmann J. The Amboseli baboon research project: 40 years of continuity and change. In: Kappeler PM, Watts DP, editors. Long-term field studies of primates. Berlin Heidelberg: Springer; 2012. p. 261–88.
- Gratten J, Pilkington JG, Brown EA, Clutton-Brock TH, Pemberton JM, Slate J. Selection and microevolution of coat pattern are cryptic in a wild population of sheep. *Mol Ecol.* 2012;21:2977–90.
- Albon SD, Coulson TN, Brown D, Guinness FE, Pemberton JM, Clutton-Brock TH. Temporal changes in key factors and key age groups influencing the population dynamics of female red deer. *J Anim Ecol.* 2000;69:1099–110.
- Bouwhuis S, Vedder O, Garroway CJ, Sheldon BC. Ecological causes of multilevel covariance between size and first-year survival in a wild bird population. *J Anim Ecol.* 2015;84:208–18.
- Grant PR, Grant BR. Unpredictable evolution in a 30-year study of Darwin's finches. *Science.* 2002;296:707–11.
- Fernandez-Duque E, Rotundo M. Field methods for capturing and marking Azarai night monkeys. *Int J Primatol.* 2003;24:1113–20.
- Van Belle S, Fernandez-Duque E, Di Fiore A. Demography and life history of wild red titi monkeys (*Callicebus discolor*) and equatorial sakis (*Pithecia aequatorialis*) in Amazonian Ecuador: A 12-year study. *Am J Primatol.* 2016;78:204–15.
- Wasserman MD, Chapman CA, Milton K, Goldberg TL, Ziegler TE. Physiological and behavioral effects of capture darting on red colobus monkeys (*Procolobus rufomitratus*) with a comparison to chimpanzee (*Pan troglodytes*) predation. *Int J Primatol.* 2013;34:1020–31.
- Wright PC. Demography and life history of free-ranging *Propithecus diadema edwardsi* in Ranomafana National Park, Madagascar. *Int J Primatol.* 1995;16:835–54.
- Richard AF, Dewar RE, Schwartz M, Ratsiraron J. Life in the slow lane? demography and life histories of male and female sifaka (*Propithecus verreauxi verreauxi*). *J Zool.* 2002;256:421–36.
- Irwin MT. Living in forest fragments reduces group cohesion in diademed sifakas (*Propithecus diadema*) in eastern Madagascar by reducing food patch size. *Am J Primatol.* 2007;69:434–47.
- Leimberger KG, Lewis RJ. Patterns of male dispersal in Verreaux's sifaka (*Propithecus verreauxi*) at Kirindy Mitea National Park. *Am J Primatol.* 2016. DOI:10.1002/ajp.22455.
- Fernandez-Duque E. Natal dispersal in monogamous owl monkeys (*Aotus azarai*) of the Argentinean Chaco. *Behaviour.* 2009;146:583–606.
- Wright PC, Erhart EM, Tecot S, Baden AL, Arrigo-Nelson SJ, Herrera J, Morelli TL, Blanco MB, Deppe A, Atsalis S, Johnson S, Ratelolahy F, Tan C, Zohdy S. Long-term research at Centre ValBio, Ranomafana National Park, Madagascar. In: Kappeler PM, Watts DP, editors. Long-term field studies of primates. Berlin Heidelberg: Springer; 2012. p. 67–100.
- Tecot S, Gerber B, King S, Verdolin J, Wright PC. Risky business: sex ratio, mortality, and group transfer in *Propithecus edwardsi* in Ranomafana National Park, Madagascar. *Behav Ecol.* 2013;24:987–96.
- Zohdy S, Gerber BD, Tecot S, Blanco MB, Winchester JM, Wright PC, Jernvall J. Teeth, sex, and testosterone: aging in the world's smallest primate. *PLoS ONE.* 2014;9:e109528.

25. Glander KE, Wright PC, Daniels PS, Merenlender AM. Morphometrics and testicle size of rain-forest lemur species from southeastern Madagascar. *J Hum Evol.* 1992;22:1–17.
26. Sorin AB. Paternity assignment for white-tailed deer (*Odocoileus virginianus*): mating across age classes and multiple paternity. *J Mammal.* 2004;85:356–62.
27. Loison A, Solberg EJ, Yoccoz NG, Langvatn R. Sex differences in the interplay of cohort and mother quality on body mass of red deer calves. *Ecology.* 2004;85:1992–2002.
28. Hopkins ME, Milton K. Adverse effects of ball-chain radio-collars on female mantled howlers (*Alouatta palliata*) in Panama. *Int J Primatol.* 2016;37:213–24.
29. Cunningham EP, Unwin S, Setchell JM. Darting primates in the field: a review of reporting trends and a survey of practices and their effect on the primates involved. *Int J Primatol.* 2015;36:911–32.
30. Côté SD, Festa-Bianchet M, Fournier F. Life-history effects of chemical immobilization and radiocollars on mountain goats. *J Wildlife Manage.* 1998; 62:745–52.
31. Moorhouse TP, Macdonald DW. Indirect negative impacts of radio-collaring: sex ratio variation in water voles. *J Appl Ecol.* 2005;42:91–8.
32. Le Maho Y, Saraux C, Durant JM, Viblanc VA, Gauthier-Clerc M, Yoccoz NG, Stenseth NC, Le Bohec C. An ethical issue in biodiversity science: the monitoring of penguins with flipper bands. *C R Biol.* 2011;334:378–84.
33. Tecot SR. It's all in the timing: out of season births and infant survival in *Eulemur rubriventer*. *Int J Primatol.* 2010;31:715–35.
34. Kühl HS, Burghardt T. Animal biometrics: quantifying and detecting phenotypic appearance. *Trends Ecol Evol.* 2013;28:432–41.
35. Kelly MJ. Computer-aided photograph matching in studies using individual identification: an example from Serengeti cheetahs. *J Mammal.* 2001;82:440–9.
36. Hiby L, Lovell P, Patil N, Kumar NS, Gopalaswamy AM, Karnath KU. A tiger cannot change its stripes: using a three-dimensional model to match images of living tigers and tiger skins. *Biol Lett.* 2009;5:383–6.
37. Bolger DT, Morrison TA, Vance B, Lee D, Farid H. A computer-assisted system for photographic mark–recapture analysis. *Methods Ecol Evol.* 2012;3:813–22.
38. Lahiri M, Tantipathananandh C, Warungu R, Rubenstein DI, Berger-Wolf TY. Biometric animal databases from field photographs: identification of individual zebra in the wild. *ICMR.* 2011. [http://compbio.cs.uic.edu/~mayank/papers/LahiriEtal_ZebraID11.pdf] Downloaded 1 July 2013.
39. Crouse D, Richardson Z, Jain A, Tecot S, Baden A, Jacobs R. Lemur face recognition: tracking a threatened species and individuals with minimal impact. MSU Technical Report 2015. MSU-CSE-15-8, May 23, 2015.
40. Li SZ, Jain AK. Handbook of face recognition. 2nd ed. London: Springer; 2011.
41. Loos A, Ernst A. An automated chimpanzee identification system using face detection and recognition. *EURASIP J Image Video Process.* 2013;1:1–17.
42. Allen AL, Higham JP. Assessing the potential information content of multicomponent visual signals: a machine learning approach. *Proc R Soc B.* 2015;282:20142284.
43. Mittermeier RA, Louis EE, Richardson M, Schwitzer C, Langrand O, Rylands AB, Hawkins F, Rajaobelina S, Ratsimbazafy J, Rasoloarison R, Roos C, Kappeler PM, MacKinnon J. Lemurs of Madagascar. Arlington: Conservation International; 2010.
44. Overdorff DJ. Ecological correlates to social structure in two prosimian primates: *Eulemur fulvus rubosus* and *Eulemur rubriventer* in Madagascar. PhD thesis. Duke University, Durham: Department of Biological Anthropology and Anatomy; 1991.
45. Durham DL. Variation in responses to forest disturbance and the risk of local extinction: a comparative study of wild *Eulemurs* at Ranomafana National Park. PhD thesis. University of California, Davis: Department of Animal Behavior; 2003.
46. Tecot SR. Seasonality and predictability: the hormonal and behavioral responses of the red-bellied lemur, *Eulemur rubriventer*, in southeastern Madagascar. PhD thesis. University of Texas at Austin, Austin: Department of Anthropology; 2008.
47. Jacobs RL. The evolution of color vision in red-bellied lemurs (*Eulemur rubriventer*). PhD thesis. Stony Brook University, Stony Brook: Department of Anthropology (Physical Anthropology); 2015.
48. Tan X, Triggs B. Enhanced local texture feature sets for face recognition under difficult lighting conditions. *IEEE T Image Process.* 2010;19:1635–50.
49. Klum S, Han H, Jain AK, Klare B. Sketch based face recognition: forensic vs. composite sketches. In *Biometrics (ICB), 2013 International Conference on Biometrics Compendium, IEEE.* 2013:1–8. DOI: 10.1109/ICB.2013.6612993.
50. Ojala T, Pietikainen M, Maenpää T. Multiresolution gray-scale and rotation invariant texture classification with local binary patterns. *IEEE T Pattern Anal.* 2002;24:971–87.
51. Lowe DG. Distinctive image features from scale-invariant keypoints. *Int J Comput Vision.* 2004;60:91–110.
52. Overdorff DJ. Similarities, differences, and seasonal patterns in the diets of *Eulemur rubriventer* and *Eulemur fulvus rufus* in the Ranomafana National Park, Madagascar. *Int J Primatol.* 1993;14:721–53.
53. Harper GJ, Steininger MK, Tucker CJ, Juhn D, Hawkins F. Fifty years of deforestation and forest fragmentation in Madagascar. *Environ Conserv.* 2007;34:325–33.
54. Andriaholinirina N, Baden A, Blanco M, Chikhi L, Cooke A, Davies N, Dolch R, Donati G, Ganzhorn J, Golden C, Groeneveld LF, Hapke A, Irwin M, Johnson S, Kappeler P, King T, Lewis R, Louis EE, Markolf M, Mass V, Mittermeier RA, Nichols R, Patel E, Rabarivola CJ, Raharivololona B, Rajaobelina S, Rakotoarisoa G, Rakotomanga B, Rakotonahary J, Rakotonrainibe H et al. *Eulemur rubriventer*. The IUCN Red List of Threatened Species 2014, Version 2015.2. Downloaded on 26 May 2016 [www.iucnredlist.org].
55. Wright PC. Primate ecology, rainforest conservation, and economic development: building a national park in Madagascar. *Evol Anthropol.* 1992;1:25–33.
56. Klontz JC, Klare BF, Klum S, Jain AK, Burge MJ. Open source biometric recognition. In *Proceedings of Biometrics: Theory, Applications and Systems (BTAS), 2013 IEEE Sixth International Conference.* 2013:1–8. [<http://openbiometrics.org/publications/klontz2013open.pdf>].
57. Yip AW, Sinha P. Contribution of color to face recognition. *Perception.* 2002; 31:995–1003.
58. Martinkauppi JB, Hadid A, Pietikainen M. Skin color in face analysis. In: Li SZ, Jain AK, editors. Handbook of face recognition. Second ed. London: Springer; 2011. p. 223–49.
59. Jain AK, Dubes RC. Algorithms for clustering data. New Jersey: Prentice Hall; 1988.
60. Otto C, Klare BF, Jain AK. An efficient approach for clustering face images. In *Proceedings of IEEE International Conference on Biometrics (ICB), Phuket, Thailand, May 19–22, 2015.* (doi: 10.1109/ICB.2015.7139091)
61. Yoder AD, Yang Z. Divergence dates for Malagasy lemurs estimated from multiple gene loci: geological and evolutionary context. *Mol Ecol.* 2004;13:757–73.
62. Crowley BE, Godfrey LR, Bankoff RJ, Perry GH, Culleton BJ, Kennett DJ, Sutherland MR, Samonds KE, Burney DA. Island-wide aridity did not trigger recent megafaunal extinctions in Madagascar. *Ecography.* 2016. DOI:10.1111/ecog.02376.
63. Brook BW, O'Grady JJ, Chapman AP, Burgman MA, Akçakaya HR, Frankham R. Predictive accuracy of population viability analysis in conservation biology. *Nature.* 2000;404:385–7.
64. Strier KB, Alberts S, Wright PC, Altmann J, Zeitley D. Primate life-history databank: setting the agenda. *Evol Anthropol.* 2006;15:44–6.
65. Strier KB, Altmann J, Brockman DK, Bronikowski AM, Cords M, Fedigan LM, Lapp H, Liu X, Morris WF, Pusey AE, Stoinski TS, Alberts SC. The Primate Life History Database: a unique shared ecological data resource. *Methods Ecol Evol.* 2010;1:199–211.
66. Reuter KE, Gilles H, Wills AR, Sewall BJ. Live capture and ownership of lemurs in Madagascar: extent and conservation implications. *Oryx.* 2016;50:344–54.
67. Reuter KE, Schaefer MS. Captive conditions of pet lemurs in Madagascar. *Folia Primatol.* 2016;2016(87):48–63.
68. Nijman V, Nekaris KAI, Donati G, Bruford M, Fa J. Primate conservation: measuring and mitigating trade in primates. *Endang Species Res.* 2011;13:159–61.
69. Bush ER, Baker SE, Macdonald DW. Global trade in exotic pets 2006–2012. *Conserv Biol.* 2014;28:663–76.
70. Jackson RM, Roe JD, Wangchuk R, Hunter DO. Estimating snow leopard population abundance using photography and capture-recapture techniques. *Wildlife Soc Bull.* 2006;34:772–81.
71. Karanth KU, Nichols JD, Kumar NS, Hines JE. Assessing tiger population dynamics using photographic capture-recapture sampling. *Ecology.* 2006;87:2925–37.
72. Jain AK, Farrokhnia F. Unsupervised texture segmentation using Gabor filters. *Pattern Recogn.* 1991;24:1167–86.
73. Freytag A, Rodner E, Simon M, Loos A, Kühl HS, Denzler J. Chimpanzee faces in the wild: Log-Euclidean CNNs for predicting identities and attributes of primates. In: Rosenhahn B, Bjoern A, editors. *Pattern recognition 38th German conference, GCPR 2016, Hannover, Germany, September 12–15, 2016, proceedings.* Switzerland: Springer International Publishing AG; 2016. p. 51–63.
74. Jacobs RL, Bradley BJ. Considering the influence of nonadaptive evolution on primate colour vision. *Plos One.* 2016;11:e0149664.