Lipids in Health and Disease

CORRECTION

Correction: The blood monocyte to high density lipoprotein cholesterol ratio (MHR) is a possible marker of carotid artery plaque

Jie Xi[1], Shasha Men[2], Jingzhu Nan[2], Qiuliang Yang[2] and Jin Dong[2*]

Correction: Lipids Health Dis 21, 130 (2022)
https://doi.org/10.1186/s12944-022-01741-8

Following publication of the original article [1], the authors reported an error in Fig. 1: The input method icon was not removed when combining the figures in Fig. 1. Figure 1 should have appeared as shown below.

The original article has been updated.

Xi *et al. Lipids in Health and Disease*   (2022) 21:144

Page 2 of 3

Fig. 1 Correlation analysis of MHR and other metabolic indexes. Abbreviations: MHR: monocyte to HDL-C ratio; BMI: body mass index; hs-CRP: high-sensitivity C-reactive protein; Glu: glucose; HbA1c: hemoglobin A1c; LDL-C: low-density lipoprotein cholesterol; TC: total cholesterol; Tg: triglyceride; Crea: creatinine; Ua: uric acid; eGFR: estimated glomerular filtration rate

Xi *et al. Lipids in Health and Disease*   (2022) 21:144

Page 3 of 3

Author details
[1]Center of Translational Medicine Research, Medical Innovation Research Department of Chinese PLA General Hospital, Beijing 100853, China. [2]Department of Clinical Laboratory, the 1st Medical Centre, Chinese PLA General Hospital, Beijing 100853, China.

Reference
1. Xi J, Men S, Nan J, et al. The blood monocyte to high density lipoprotein cholesterol ratio (MHR) is a possible marker of carotid artery plaque. Lipids Health Dis. 2022;21:130. https://doi.org/10.1186/s12944-022-01741-8.