## CORRECTION Open Access

## Correction: The relationship between Lipocalin-2 level and hepatic steatosis in obese patients with NAFLD after bariatric surgery

Jiaqi Chen<sup>1,2</sup>, Shihui Lei<sup>1</sup>, Yueye Huang<sup>1</sup>, Xiaojuan Zha<sup>1</sup>, Lei Gu<sup>3</sup>, Donglei Zhou<sup>3</sup>, Jun Li<sup>4</sup>, Feng Liu<sup>4</sup>, Nannan Li<sup>1</sup>, Lei Du<sup>1</sup>, Xiu Huang<sup>1</sup>, Ziwei Lin<sup>1</sup>, Le Bu<sup>1\*</sup> and Shen Qu<sup>1\*</sup>

## Correction: Lipids Health Dis 21, 10 (2022) https://doi.org/10.1186/s12944-022-01622-0

Following publication of the original article [1], the authors reported an error in affiliations.

The accurate author affiliations should be as follows:

1. Department of Endocrinology and Metabolism, Shanghai Tenth People's Hospital, Clinical Medicine School of Nanjing Medical University, Medicine School of Tongji University, Shanghai 200072, China.

- 2. Department of Endocrinology and Metabolism, Suzhou Municipal Hospital, The Affiliated Suzhou Hospital of Nanjing Medical University, Suzhou, China.
- 3. Department of Gastrointestinal Surgery, Shanghai Tenth People's Hospital, Clinical Medicine School of Nanjing Medical University, Medicine School of Tongji University, Shanghai 200072, China.
- 4. Department of Gastroenterology, Shanghai Tenth People's Hospital, Clinical Medicine School of Nanjing Medical University, Medicine School of Tongji University, Shanghai 200072, China.

The original article has been updated.

The original article can be found online at https://doi.org/10.1186/s12944-022-01622-0.

\*Correspondence:

Le Bu

geyingjun@hotmail.com

Shen Qu

qushencn@hotmail.com

- <sup>1</sup> Department of Endocrinology and Metabolism, Shanghai Tenth People's Hospital, Clinical Medicine School of Nanjing Medical University, Medicine School of Tongji University, Shanghai 200072, China
- <sup>2</sup> Department of Endocrinology and Metabolism, Suzhou Municipal Hospital, The Affiliated Suzhou Hospital of Nanjing Medical University, Suzhou, China
- <sup>3</sup> Department of Gastrointestinal Surgery, Shanghai Tenth People's Hospital, Clinical Medicine School of Nanjing Medical University, Medicine School of Tongji University, Shanghai 200072, China
- <sup>4</sup> Department of Gastroenterology, Shanghai Tenth People's Hospital, Clinical Medicine School of Nanjing Medical University, Medicine School of Tongji University, Shanghai 200072, China

Published online: 27 February 2023

## Reference

 Chen J, Lei S, Huang Y, et al. The relationship between Lipocalin-2 level and hepatic steatosis in obese patients with NAFLD after bariatric surgery. Lipids Health Dis. 2022;21:10. https://doi.org/10.1186/ s12944-022-01622-0.



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/joublicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.