

BRIEF REPORT

Open Access



A pilot study of the association between maternal mid-pregnancy cholesterol and oxysterol concentrations and labor duration

Todd C. Rideout^{1*}, Jaclyn Wallace², Xiaozhong Wen³, Vanessa M. Barnabei⁴, Kai Ling Kong^{5,6,7} and Richard W. Browne²

Abstract

Background Previous animal model studies have highlighted a role for cholesterol and its oxidized derivatives (oxysterols) in uterine contractile activity, however, a lipotoxic state associated with hypercholesterolemia may contribute to labor dystocia. Therefore, we investigated if maternal mid-pregnancy cholesterol and oxysterol concentrations were associated with labor duration in a human pregnancy cohort.

Methods We conducted a secondary analysis of serum samples and birth outcome data from healthy pregnant women ($N = 25$) with mid-pregnancy fasting serum samples collected at 22–28 weeks of gestation. Serum was analyzed for total-C, HDL-C, and LDL-C by direct automated enzymatic assay and oxysterol profile including 7 α -hydroxycholesterol (7 α OHC), 7 β -hydroxycholesterol (7 β OHC), 24-hydroxycholesterol (24OHC), 25-hydroxycholesterol (25OHC), 27-hydroxycholesterol (27OHC), and 7-ketocholesterol (7KC) by liquid chromatography-selected ion monitoring-stable isotope dilution-atmospheric pressure chemical ionization-mass spectroscopy. Associations between maternal second trimester lipids and labor duration (minutes) were assessed using multivariable linear regression adjusting for maternal nulliparity and age.

Results An increase in labor duration was observed for every 1-unit increment in serum 24OHC (0.96 min [0.36,1.56], $p < 0.01$), 25OHC (7.02 min [1.92,12.24], $p = 0.01$), 27OHC (0.54 min [0.06, 1.08], $p < 0.05$), 7KC (8.04 min [2.7,13.5], $p < 0.01$), and total oxysterols (0.42 min [0.18,0.06], $p < 0.01$). No significant associations between labor duration and serum total-C, LDL-C, or HDL-C were observed.

Conclusions In this cohort, mid-pregnancy concentrations of maternal oxysterols (24OHC, 25OHC, 27OHC, and 7KC) were positively associated with labor duration. Given the small population and use of self-reported labor duration, subsequent studies are required for confirmation.

Keywords Cholesterol, Oxysterols, Labor dystocia, Labor duration, Pregnancy

*Correspondence:

Todd C. Rideout

rideout@buffalo.edu

Full list of author information is available at the end of the article



This is a U.S. Government work and not under copyright protection in the US; foreign copyright protection may apply 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

An increase in maternal lipids during pregnancy is recognized as a normal physiological response to pregnancy, with large increases in triglycerides (~200–400%), total-cholesterol (~25–50%), and low-density lipoprotein (LDL-C, ~70%) beginning in early pregnancy and peaking in the third trimester [1, 2]. However, recent work has highlighted concerns that an abnormal dyslipidemic profile during pregnancy may have negative health implications for both the mother and fetus [3, 4]. We previously reported that total particle concentrations of very low-density lipoprotein/chylomicron remnants (VLDL/CM) and high-density lipoproteins (HDL) in mid-pregnancy have divergent associations with birth weight [5]. In this brief report, we extend our focus to examine the association between maternal pregnancy cholesterol and oxysterol concentrations and labor duration.

Labor dystocia, defined as an abnormally slow or protracted labor [6], increases the risk of both maternal and neonatal complications [7, 8] and is a primary indication for cesarean delivery [6]. Predictive maternal biomarkers of a labor dystocia phenotype may be useful in identifying at-risk pregnancies and initiating early monitoring and prevention strategies. A potential link between lipotoxicity and labor dystocia has been suggested by previous human and animal model studies demonstrating a critical role for cholesterol in successful parturition but impaired uterine function and contractile activity in hypercholesterolemia states [9–12].

Several regulatory proteins controlling myometrial contraction, including the oxytocin receptor (OXTR) and human ether-a-go-go-related gene (hERG) potassium channel proteins [13] are localized in cholesterol-rich microdomains (lipids rafts and caveolae) and are therefore sensitive to local and circulating cholesterol concentrations [10, 14]. The OXTR expresses a cholesterol binding site which is important for receptor stabilization and posttranslational processing [15–17]. Further, oxysterols, a class of highly reactive oxygenated derivatives of cholesterol that induce inflammation have been shown to function as nuclear receptor ligands to regulate intracellular cholesterol concentration in the mouse uterus [18]. However, despite the known function of cholesterol and its oxidative derivatives in regulating uterine contractile activity, we are not aware of any previous human cohorts that have examined if associations exist between maternal sterol status during pregnancy and labor duration. Therefore, in this pilot study, we sought to examine if maternal mid-pregnancy cholesterol and oxysterol concentrations were associated with labor duration.

Methods

Participants

We undertook a secondary analysis of serum samples and birth outcome data collected as part of a previous pregnancy pilot cohort study. All procedures were approved by the Institutional Review Board at the University at Buffalo (STUDY00001381). Details on the study design and cohort have previously been published [5]. Briefly, the cohort consisted of 25 adult pregnant women aged 18–35 years with singleton gestations recruited within the greater Buffalo, New York area between 2018 and 2019 through flyer distribution, social media posts, and local urban obstetrics clinics (Table 1). Women were excluded if they had a history of major chronic diseases (e.g., heart disease, diabetes) or were currently taking medications that affected appetite or insulin sensitivity. None of the mothers included in this cohort had premature infants or cesarean deliveries.

Data collection

At enrollment, prospective women completed an interviewer-administered questionnaire to obtain information regarding general socio-demographics as well as reproductive and medical history. Maternal height and pre-pregnancy weight were collected by self-report at the initial visit. Body mass index (BMI) was calculated as pre-pregnancy weight in kilograms divided by the square of height in meters. At mid-pregnancy, between gestation weeks 22–28, body weight and height were directly measured by research staff using a digital scale and digital stadiometer, respectively. At this same visit, a fasting blood sample was collected by a blood draw technician. The total duration of active labor (>4 cm dilation) was obtained with a self-reported birth outcome questionnaire assessed at a home visit within 7 days of birth.

Serum lipid analyses

Following a blood clotting period of 30 minutes at room temperature, serum samples were separated by centrifugation at 3000 x g for 20 min and stored at –80 °C. Total-cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), and high-density lipoprotein cholesterol (HDL-C) were measured by direct automated enzymatic assay using diagnostic reagent kits, calibrators, and quality controls.

Serum oxysterols including 7 α -hydroxycholesterol (7 α OHC), 7 β -hydroxycholesterol (7 β OHC), 24-hydroxycholesterol (24OHC), 25-hydroxycholesterol (25OHC), 27-hydroxycholesterol (27OHC), and 7-ketocholesterol (7KC) were determined by liquid chromatography-selected ion monitoring-stable isotope dilution-atmospheric pressure chemical ionization-mass spectroscopy

Table 1 Characteristics of mothers in the pilot cohort

Variable (N = 25)	Mean (SD)	Median (IQ)	n (%)	Range
Age (yrs)	30.7 (3.96)			21.6–37.8
Race				
White			23 (92)	
African American			1 (4.0)	
Asian			1 (4)	
Education				
Level (yrs)	16.1 (2.1)			
College graduate			23 (92)	
Pre-Pregnancy BMI (kg/m²)^a				
Normal			10 (40)	
Overweight			7 (28)	
Obese			8 (32)	
Nulliparous			16 (64)	
Length of active labor (hrs)		7.0 (9.0)		0.67–30
Labor				
Labor induction			8 (32)	
Use of pain medication ^b			19 (76)	
Gestational age^c		39 (1.0)		32–41
Birth weight (g)	3543 (333)			
Mid-Pregnancy Serum Lipids (mg/dL)				
Total-C	250.6 (49.3)			131–374
LDL-C	141.7 (41.1)			40–250
HDL-C	78.9 (20.8)			33–117
Triglycerides	172.8 (61.4)			87–398
Mid-Pregnancy Serum Oxysterols (ng/mL)^d				
24-hydroxycholesterol (24OHC)	84.9 (29.3)			(42.8–146.9)
25-hydroxycholesterol (25OHC)	18.4 (3.5)			(10.7–27.0)
27-hydroxycholesterol (27OHC)	144.5 (39.3)			(66.9–211.7)
7 α -hydroxycholesterol (7 α OHC)	47.4 (27.0)			(13.4–122.2)
7 β -hydroxycholesterol (7 β OHC)	7.6 (5.0)			(0.06–21.3)
7-ketocholesterol (7KC)	5.0 (2.9)			(1.2–12.3)
Total oxysterols ^a	307.2 (72.0)			(145.9–473.9)

^a Normal BMI 18.5–24.9; overweight BMI 25–29.9; obese BMI 30–34.9

^b Including epidural analgesia (n = 16) and narcotic pain medications (n = 3)

^c Gestational weeks at time of birth

^d Represents the sum of 24OHC, 25OHC, 27OHC, 7 α OHC, 7 β OHC, and 7KC

(LC-SIM-SID-APCI-MS) as previously described [19, 20] with slight modification. Briefly, serum oxysterols were released from steryl esters by incubation with microbial cholesterol esterase (100ul containing 2U for 20 minutes at 37°C) and following solid phase extraction, were injected into an Ascentis 10 cm X 3 mm, 3- μ m C-18 LC column (SupelCo Inc.) on a Shimadzu 10ADVP LC system interfaced with a Shimadzu 2010A MS by APCI interface. Deuterated (d7) 22HC, (d7) 7 α HC and (d7) 7KC was used as stable, isotope-labeled, internal standards. We, and others [21], have found that oxysterol analytes are stable at -80°C storage. The assay was previously

validated in accordance with the FDA guidance for bio-analytical methods [22].

Statistical analyses

The relationship of individuals oxysterols with serum cholesterol and maternal BMI were assessed with Pearson correlation coefficients. Associations between maternal second trimester lipids and labor duration were assessed using multivariable linear regression. As labor duration data was not normally distributed, the variable was log transformed for all analyses. We considered nulliparity (yes/no), birth weight, gestation age, maternal age

and pre-pregnancy BMI (continuous variables) as potential confounding factors. Confounders with a (p -value < 0.05) were considered significant and were further adjusted in the models. Two models were used in the analysis: (i) Model 1 was the crude model with second trimester lipid exposure only; and (ii) Model 2 adjusted for maternal nulliparity and maternal age (the only two significant confounders). All analyses were conducted using SAS version 9.4 (SAS Institute, Inc., Cary, NC). Statistical significance was set at a two-sided alpha level of 0.05. A post hoc power analysis, based on linear bivariate regression between maternal total oxysterols and labor duration, with significance set at $\alpha = 0.05$, yielded a power ($1 - \beta$) of 0.766 (G*Power 3.1.5) [23].

Results

Mothers included in this study were mainly White (92%), highly educated (92% college graduate), and overweight (28%) or obese (32%) before pregnancy (Table 1). The majority of women 16 (64%) in the cohort were nulliparous. Mean active labor duration was 8.9 ± 6.9 hours (13.9 ± 8.6 hours for nulliparous women; 6.2 ± 4.2 hours multiparous women).

Serum oxysterols were not correlated ($p > 0.05$) with maternal BMI at pre-pregnancy or mid-pregnancy (Table 2). Serum 24OHC ($r = 0.37$), 27OHC ($r = 0.49$), 7KC ($r = 0.33$), and total oxysterols ($r = 0.52$) were positively correlated with total-C, however, no correlation ($p > 0.05$) was observed between cholesterol and 25OHC, 7 α OHC, and 7 β OHC. Amongst the oxysterols, only 27OHC and total oxysterols had significant ($p < 0.05$) positive associations with LDL-C ($r = 0.43$ and 0.41) and HDL-C ($r = 0.51$ and 0.42).

No significant associations between labor duration and serum total-C ($p = 0.08$), LDL-C ($p = 0.07$), and HDL-C ($p = 0.93$) were observed (Table 3). Among the oxysterols, with the exception of 7 α OHC and 7 β OHC, an increase in labor duration was observed for every 1-unit increment in serum serum 24OHC (0.96 min [0.36,1.56], $p < 0.01$), 25OHC (7.02 min [1.92,12.24], $p = 0.01$), 27OHC (0.54 min [0.06, 1.08], $p < 0.05$), 7KC (8.04 min [2.7,13.5], $p < 0.01$), and total oxysterols (0.42 min [0.18,0.06], $p < 0.01$) (Table 3).

Discussion

In this cohort, maternal mid-pregnancy serum oxysterol concentrations (24OHC, 25OHC, 27OHC, and 7KC) were positively associated with labor duration. However, the magnitude of these associations varied depending on the oxysterol species, possibly reflecting their specific routes of synthesis and functional roles. Interestingly, serum TC and LDL-C were not associated with labor duration. This contrast is perhaps surprising given the significant correlations we observed between serum total oxysterols and cholesterol and the known role of oxysterols in regulating intracellular cholesterol efflux and accumulation [18]. Alternatively, due to the specific roles of oxysterols as inflammatory mediators and nuclear receptor ligands, oxysterols may represent a more sensitive marker of uterine contraction than cholesterol.

Oxysterols are highly reactive oxygenated derivatives of cholesterol that differ in their oxidation sites (within the A or B rings or the side chain of cholesterol) and their routes of synthesis [24]. They can be generated through specific cytochrome P450 mitochondrial or microsomal enzymes that are coordinately regulated through

Table 2 Correlations between maternal mid-pregnancy oxysterol concentrations, weight status and serum lipids

Maternal characteristic ($n = 25$)		Mid-Pregnancy Serum Oxysterols (ng/mL)						
		24OHC	25OHC	27OHC	7 α OHC	7 β OHC	7 KC	Total Oxysterols ^a
Weight status								
Pre-pregnancy BMI (kg/m ²)	r	-0.19	-0.13	-0.21	-0.02	-0.04	0.00	-0.20
	p	0.31	0.31	0.27	0.94	0.81	1.00	0.27
Mid-pregnancy BMI (kg/m ²)	r	-0.15	-0.11	-0.15	0.03	-0.08	0.05	-0.14
	p	0.42	0.42	0.41	0.88	0.66	0.80	0.44
Serum Lipids								
Total-C (mg/dL)	r	0.37	0.19	0.49	0.23	-0.15	0.33	0.52
	p	0.04	0.30	0.004	0.21	0.41	0.07	0.002
LDL-C (mg/dL)	r	0.28	0.20	0.43	0.14	-0.22	0.21	0.41
	p	0.12	0.26	0.01	0.45	0.23	0.25	0.02
HDL-C (mg/dL)	r	0.22	0.10	0.51	0.10	-0.10	0.32	0.42
	p	0.23	0.60	0.00	0.57	0.60	0.07	0.02

r, Pearson Correlation Coefficient; p , p -value; ^aRepresents the sum of 24OHC, 25OHC, 27OHC, 7 α OHC, 7 β OHC, and 7KC

Table 3 Associations between maternal oxysterol concentrations and labor duration

Serum lipids	Labor Duration (minutes)			
	Model 1 β (95% CI) N = 25	p-value	Model 2 β (95% CI) N = 25	p-value
Lipid panel (mg/dL)				
Total-cholesterol	0.36 (−0.18,1.02)	0.22	0.48 (−0.06,0.96)	0.08
LDL-cholesterol	0.48 (−0.36,1.32)	0.25	0.60 (−0.06,1.32)	0.07
HDL-cholesterol	0.06 (−1.2,1.26)	0.97	−0.06(−1.02,0.96)	0.93
Oxysterols (ng/mL)				
24-hydroxycholesterol	1.14 (0.42,1.86)	<0.01	0.96 (0.36,1.56)	<0.01
25-hydroxycholesterol	6.9 (0.24,13.62)	0.04	7.02 (1.92,12.24)	0.01
27-hydroxycholesterol	0.72 (0.06,1.32)	0.03	0.54 (0.06,1.08)	0.04
7α-hydroxycholesterol	0.24 (−0.48,1.08)	0.50	0.42 (−0.18,1.02)	0.20
7β-hydroxycholesterol	1.56 (−3.36,6.48)	0.53	1.02 (−3.06,5.1)	0.62
7-ketocholesterol	7.38 (0.42,14.34)	0.03	8.04 (2.7,13.5)	<0.01
Total oxysterols	0.48 (0.24,0.72)	<0.01	0.42 (0.18,0.06)	<0.01

*Values are beta coefficients (95% CI) derived from linear regression models, representing the change in labor duration (minutes) with one unit increment in each lipid exposure; Model 1: crude model; Model 2: adjusted for maternal nulliparity and age

substrate and co-factor availability or through random non-enzymatic autooxidation routes involving reactive oxygen species [25]. Oxysterols of enzymatic origin, including 7αOHC, 24OHC, 25OHC, and 27OHC, are most often identified as intermediates of bile acid synthesis, however, they also act as intracellular ligands for several nuclear receptors and thus play a functional role in regulating a broad range of metabolic pathways, including cholesterol metabolism. Alternatively, oxysterols of non-enzymatic origin, including 7βOHC and 7KC, are typically recognized for their pro-oxidant and pro-inflammatory biological activities through their induction of inflammatory cytokine expression [24].

Although oxysterols have been implicated in the pathogenesis of several disease states including atherosclerosis [26] and dementia [27], few studies have characterized oxysterol concentrations during pregnancy or examined their relationship with pregnancy outcomes. In a longitudinal pregnancy cohort of 33 women, Winkler et al. (2017) [28] reported an increase in maternal 27OHC throughout pregnancy but no association with several pregnancy complications including intrauterine growth restriction (IUGR), preeclampsia (PE), and hemolysis, elevated liver enzymes, low platelet count (HELLP) syndrome. However, Mistry et al. observed an increase in placental 27OHC concentration and enhanced maternal and fetal cholesterol-mediated serum efflux capacity in pregnancies affected by preeclampsia compared with normotensive controls [29]. A potential role of oxysterols in placental dysfunction is further supported through in vitro studies reporting the pro-inflammatory actions of

25OHC and 7KC through a toll-like receptor 4 (TLR4)-dependent mechanism [30] and cytotoxic effects of 25OHC in placental trophoblasts [31].

We are not aware of previous human studies that have examined the association between maternal cholesterol or their oxygenated derivatives and labor duration. The observed differences in the relative magnitude of the effect for individual oxysterol species may be of interest in their potential utility as predictive biomarkers of labor dystocia. At this point, the underlying reasons for these differential responses are not known. However, while the physiological effects of oxysterols as a class of compounds has been historically emphasized [32], more recent work has recognized the metabolic implications of individual oxysterol species which may be influenced by several factors including absolute concentrations and tissue distribution [33], differential ligand binding affinities [34], and a diverse range of metabolic activities [24].

Although our findings cannot provide causative evidence to support a direct role of maternal sterol status on labor duration, results from animal model studies suggest several mechanistic links between lipotoxicity (specifically, cholesterol and oxysterols) and labor dystocia. First, as the precursor substrate of steroid hormones, cholesterol is essential for the endocrine balance between estrogen and progesterone that activates the myometrium and regulates the onset of labor [35]. Second, under normal physiological concentrations, cholesterol is involved in the stabilization and posttranslational processing of the oxytocin receptor [15, 16], however, an abnormal increase in cholesterol

has been shown to blunt oxytocin-induced contractile activity in mouse [12] and human [11] uterine tissue. Third, previous work by Mouzat et al. (2006) in mice suggests that oxysterols may have a role in preserving uterine contractile function under excessive cholesterol conditions by signaling increased transcription of cholesterol efflux genes through ligand activation of the liver X receptor β (LXR β) [18].

The results of this pilot study should be interpreted with caution given several study limitations. First, due to resource constraints, this study included 25 pregnant women of which only 64% were nulliparous and for whom labor dystocia is more commonly diagnosed. However, we hope the results provide insight into estimated effect size for the design of future randomized controlled trials to identify lipid biomarkers of labor dystocia. Second, our sample was limited with respect to race/ethnicity and education and may be biased due to potentially inaccurate assessment of labor duration with a self-reported birth outcome questionnaire. Finally, oxysterol analysis was limited to mid-pregnancy only, and it is unknown how concentrations at this timepoint correlate with those at early and late gestation phases.

Conclusions

In this cohort, mid-pregnancy concentrations oxysterols (24OHC, 25OHC, 27OHC, and 7KC) were positively associated with labor duration. However, considering the small population and use of self-reported labor duration, subsequent studies are required to support the potential use of serum oxysterols as biomarkers of labor duration.

Abbreviations

OXR	Oxytocin receptor
hERG	Human ether-a-go-go-related gene
BMI	Body mass index
TC	Total-cholesterol
LDL-C	Low-density lipoprotein cholesterol
HDL-C	High-density lipoprotein cholesterol
7 α OHC	7 α -hydroxycholesterol
7 β OHC	7 β -hydroxycholesterol
24OHC	24-hydroxycholesterol
25OHC	25-hydroxycholesterol
27OHC	27-hydroxycholesterol
7KC	7-ketocholesterol
LC-SIM-SID-APCI-MS	Liquid chromatography-selected ion monitoring-stable isotope dilution-atmospheric pressure chemical ionization-mass spectroscopy
IUGR	Intrauterine growth restriction
PE	Preeclampsia
HELLP	Hemolysis, elevated liver enzymes, low platelet count syndrome
TLR4	Toll-like receptor 4
LXR β	Liver X receptor β

Acknowledgements

We are grateful to the Clinical and Translational Science Institute at the University at Buffalo for their help in supporting this work.

Authors' contributions

TCR, XW, VB, KLK, and RWB conceived the study and contributed to data interpretation and manuscript preparation. RWB and JW conducted the oxysterol analysis. TCR analyzed the data and wrote the initial draft manuscript. All authors read, revised, and approved the final manuscript.

Funding

Research reported in this publication was supported by the National Center for Advancing Translational Sciences of the National Institutes of Health under award number UL1TR001412 to the University at Buffalo. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

All methods and procedures were carried out in accordance with relevant guidelines and regulations and approved by the Institutional Review Board at the University at Buffalo (STUDY00001381). Informed consent was obtained from all participants.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹Department of Exercise and Nutrition Sciences, School of Public Health and Health Professions, Jacobs School of Medicine and Biomedical Sciences, University at Buffalo, Buffalo, NY 14214, USA. ²Department of Biotechnical and Clinical Laboratory Sciences, Jacobs School of Medicine and Biomedical Sciences, University at Buffalo, Buffalo, NY 14214, USA. ³Department of Pediatrics, Jacobs School of Medicine and Biomedical Sciences, University at Buffalo, Buffalo, NY 14214, USA. ⁴Department of Obstetrics and Gynecology, Jacobs School of Medicine and Biomedical Sciences, University at Buffalo, Buffalo, NY 14214, USA. ⁵Baby Health Behavior Lab, Division of Health Services and Outcomes Research, Children's Mercy Research Institute, Children's Mercy Hospital, Kansas City, USA. ⁶Department of Pediatrics, University of Missouri, Columbia, USA. ⁷Center for Children's Healthy Lifestyles and Nutrition, University of Kansas Medical Center, Kansas City, KS, USA.

Received: 10 January 2023 Accepted: 3 March 2023

Published online: 11 March 2023

References

- Wang J, Li Z, Lin L. Maternal lipid profiles in women with and without gestational diabetes mellitus. *Medicine (Baltimore)*. 2019;98(16):e15320.
- Wiznitzer A, Mayer A, Novack V, Sheiner E, Gilutz H, Malhotra A, et al. Association of lipid levels during gestation with preeclampsia and gestational diabetes mellitus: a population-based study. *Am J Obstet Gynecol*. 2009;201(5):482 e1–8.
- Zhu SM, Zhang HQ, Li C, Zhang C, Yu JL, Wu YT, et al. Maternal lipid profile during early pregnancy and birth weight: a retrospective study. *Front Endocrinol (Lausanne)*. 2022;13:951871.
- Xi F, Chen H, Chen Q, Chen D, Chen Y, Sagnelli M, et al. Second-trimester and third-trimester maternal lipid profiles significantly correlated to LGA and macrosomia. *Arch Gynecol Obstet*. 2021;304(4):885–94.
- Rideout TC, Wen X, Choudhary D, Catanzaro M, Browne RW, Barnabei VM, et al. Associations of maternal lipoprotein particle distribution in mid-pregnancy with birth outcomes: a pilot study. *Lipids Health Dis*. 2022;21(1):53.
- LeFevre NM, Krumm E, Cobb WJ. Labor Dystocia in nulliparous women. *Am Fam Physician*. 2021;103(2):90–6.

7. Myers ER, Sanders GD, Coeytaux RR, et al. Labor Dystocia. Rockville (MD): Agency for Healthcare Research and Quality (US); 2020. (Comparative Effectiveness Review, No. 226).
8. Niemczyk NA, Ren D, Stapleton SR. Associations between prolonged second stage of labor and maternal and neonatal outcomes in freestanding birth centers: a retrospective analysis. *BMC Pregnancy Childbirth*. 2022;22(1):99.
9. Carlsson NS, Frediani JK, Corwin EJ, Dunlop A, Jones D. Metabolomic pathways predicting labor Dystocia by maternal body mass index. *AJP Rep*. 2020;10(1):e68–77.
10. Smith RD, Babychuk EB, Noble K, Draeger A, Wray S. Increased cholesterol decreases uterine activity: functional effects of cholesterol alteration in pregnant rat myometrium. *Am J Physiol Cell Physiol*. 2005;288(5):C982–8.
11. Jie Z, Kendrick A, Quenby S, Wray S. Contractility and calcium signaling of human myometrium are profoundly affected by cholesterol manipulation: implications for labor? *Reprod Sci*. 2007;14(5):456–66.
12. Padol AR, Sukumaran SV, Sadam A, Kesavan M, Arunvikram K, Verma AD, et al. Hypercholesterolemia impairs oxytocin-induced uterine contractility in late pregnant mouse. *Reproduction*. 2017;153(5):565–76.
13. Parkington HC, Stevenson J, Tonta MA, Paul J, Butler T, Maiti K, et al. Diminished hERG K⁺ channel activity facilitates strong human labour contractions but is dysregulated in obese women. *Nat Commun*. 2014;5:4108.
14. Balijepalli RC, Delisle BP, Balijepalli SY, Foell JD, Slind JK, Kamp TJ, et al. Kv11.1 (ERG1) K⁺ channels localize in cholesterol and sphingolipid enriched membranes and are modulated by membrane cholesterol. *Channels (Austin)*. 2007;1(4):263–72.
15. Gimpl G, Fahrenholz F. Cholesterol as stabilizer of the oxytocin receptor. *Biochim Biophys Acta*. 2002;1564(2):384–92.
16. Muth S, Fries A, Gimpl G. Cholesterol-induced conformational changes in the oxytocin receptor. *Biochem J*. 2011;437(3):541–53.
17. Wiegand V, Gimpl G. Specification of the cholesterol interaction with the oxytocin receptor using a chimeric receptor approach. *Eur J Pharmacol*. 2012;676(1–3):12–9.
18. Mouzat K, Prod'Homme M, Volle DH, Sion B, Dechelotte P, Gauthier K, et al. Oxysterol nuclear receptor LXRbeta regulates cholesterol homeostasis and contractile function in mouse uterus. *J Biol Chem*. 2007;282(7):4693–701.
19. Narayanaswamy R, Iyer V, Khare P, Bodziak ML, Badgett D, Zivadinov R, et al. Simultaneous determination of oxysterols, cholesterol and 25-hydroxy-vitamin D3 in human plasma by LC-UV-MS. *PLoS One*. 2015;10(4):e0123771.
20. Fellows Maxwell K, Bhattacharya S, Bodziak ML, Jakimovski D, Hagemeyer J, Browne RW, et al. Oxysterols and apolipoproteins in multiple sclerosis: a 5 year follow-up study. *J Lipid Res*. 2019;60(7):1190–8.
21. Helmschrodt C, Becker S, Thiery J, Ceglarek U. Preanalytical standardization for reactive oxygen species derived oxysterol analysis in human plasma by liquid chromatography-tandem mass spectrometry. *Biochem Biophys Res Commun*. 2014;446(3):726–30.
22. Food US, Administration D. FDA guidance for industry: bioanalytical method validation. In: US Department of Health and Human Services, Food and Drug Administration. Rockville: Center for Drug Evaluation and Research; 2001.
23. Faul F, Erdfelder E, Lang AG, Buchner A. G*power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods*. 2007;39(2):175–91.
24. Poli G, Biasi F, Leonarduzzi G. Oxysterols in the pathogenesis of major chronic diseases. *Redox Biol*. 2013;1:125–30.
25. Griffiths WJ, Wang Y. Oxysterols as lipid mediators: their biosynthetic genes, enzymes and metabolites. *Prostaglandins Other Lipid Mediat*. 2020;147:106381.
26. Ravi S, Duraisamy P, Krishnan M, Martin LC, Manikandan B, Raman T, et al. An insight on 7- ketocholesterol mediated inflammation in atherosclerosis and potential therapeutics. *Steroids*. 2021;172:108854.
27. Staurengi E, Cerrato V, Gamba P, Testa G, Giannelli S, Leoni V, et al. Oxysterols present in Alzheimer's disease brain induce synaptotoxicity by activating astrocytes: a major role for lipocalin-2. *Redox Biol*. 2021;39:101837.
28. Winkler BS, Pecks U, Najjari L, Kleine-Eggebrecht N, Maass N, Mohaupt M, et al. Maternal 27-hydroxycholesterol concentrations during the course of pregnancy and in pregnancy pathologies. *BMC Pregnancy Childbirth*. 2017;17(1):106.
29. Mistry HD, Kurlak LO, Mansour YT, Zurkinden L, Mohaupt MG, Escher G. Increased maternal and fetal cholesterol efflux capacity and placental CYP27A1 expression in preeclampsia. *J Lipid Res*. 2017;58(6):1186–95.
30. Aye IL, Waddell BJ, Mark PJ, Keelan JA. Oxysterols exert proinflammatory effects in placental trophoblasts via TLR4-dependent, cholesterol-sensitive activation of NF-kappaB. *Mol Hum Reprod*. 2012;18(7):341–53.
31. Larkin JC, Sears SB, Sadovsky Y. The influence of ligand-activated LXR on primary human trophoblasts. *Placenta*. 2014;35(11):919–24.
32. Schroeffer GJ Jr. Oxysterols: modulators of cholesterol metabolism and other processes. *Physiol Rev*. 2000;80(1):361–554.
33. Yutuc E, Angelini R, Baumert M, Mast N, Pikuleva I, Newton J, et al. Localization of sterols and oxysterols in mouse brain reveals distinct spatial cholesterol metabolism. *Proc Natl Acad Sci U S A*. 2020;117(11):5749–60.
34. Wang Y, Kumar N, Solt LA, Richardson TI, Helvering LM, Crumbley C, et al. Modulation of retinoic acid receptor-related orphan receptor alpha and gamma activity by 7-oxygenated sterol ligands. *J Biol Chem*. 2010;285(7):5013–25.
35. Kota SK, Gayatri K, Jammula S, Kota SK, Krishna SV, Meher LK, et al. Endocrinology of parturition. *Indian J Endocrinol Metab*. 2013;17(1):50–9.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

