

LETTER TO THE EDITOR

Open Access



# Vitamin D levels and COVID 19 risk and death; is there an association?

Maisa Hamed Al Kiyumi\*

Dear editor,

I read with great interest the study entitled “Low vitamin D levels do not aggravate COVID-19 risk or death, and vitamin D supplementation does not improve outcomes in hospitalized patients with COVID-19: a meta-analysis and GRADE assessment of cohort studies and RCTs” by Chen et al. [1]. This meta-analysis refuted any association between low vitamin D level and COVID-19 risk or death. However, I have the following comments related to the internal validity of this study:

1- The pooled analysis in Figure 2 revealed a nonsignificant association between vitamin D level and COVID-19 risk and death. However, it is obvious that the pooled effect size might be influenced by a study conducted by Hastie et al. [2, 3]. This study was based on participants recruited from the UK Biobank and did not show any association between low vitamin D level and the risk of developing COVID-19 after adjustment for potential confounders [2, 3]. Notably, the true prevalence of COVID-19 might be underestimated in this cohort because the PCR test results were available for only 1,474 participants out of 348,598 recruited [2]. Moreover, we need to interpret the findings of this study cautiously, as the baseline vitamin D level measurement was done more than a decade ago (2006–2020) [2]. Similarly, in another study by Katz et al. vitamin D measurements were performed during the preceding five years prior to the COVID 19 test [4]. Besides, it was not known

whether those recruited participants with low vitamin D level received a treatment or not, which eventually might affect the accuracy of the data. I believe that carrying out sensitivity analysis in this case might help in mitigating the influence of these studies on the overall effect size.

2- Confounding factors and methods of measuring vitamin D level: While using the adjusted odds ratio plays a key role in attenuating the effects of the confounding variables, the studies included differ considerably on the types of confounding adjustment. For example, the included odds ratio by Kanz et al. [4] were adjusted for obesity only (OR=2.27; 95% CI, 1.787–2.872;  $P < 0.001$ ), whereas Kaufmann had adjusted for race/ethnicity, gender and latitude [5]. Also, with the exception of one study by Meltzer et al. [6], the possibility of vitamin D supplement initiation or adjustment after testing was not addressed and therefore might affect the findings of this meta-analysis. Importantly, the authors did not mention the assay used for measuring vitamin D level, which might also have impacted the accuracy of the overall findings. For instance, a study substantiates a higher concentration of vitamin D level when measured by a liquid chromatography-mass spectrometry technique (LC-MS/MS) compared to radioimmunoassay, with a mean difference of about 12.9 ng/ml [7].

3- In the statistical analysis section, the authors carried out the pooled analysis using a random effect model. In view of the few studies included in the quantita-

\*Correspondence: maysa8172@gmail.com

Department of Family Medicine and Public Health, Sultan Qaboos University Hospital, P.C.123, Al-Khod, P.O.Box 35, Muscat, Sultanate of Oman



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

tive analysis, the choice of which effect model to use (random versus fixed effects model) needs to be explained to the reader. While the random effects model can still be applied, the fixed effects model might be more suitable in this meta-analysis [8].

Considering the above factors, the results of this analysis need to be cautiously interpreted, and well-designed prospective clinical trials are necessary.

#### Acknowledgements

Nil

#### Author's contributions

Conceptualization, writing and editing the manuscript. The author(s) read and approved the final manuscript.

#### Funding

This manuscript is not funded.

#### Availability of data and materials

Not applicable.

#### Declarations

#### Ethics approval and consent to participate

Not applicable.

#### Consent for publication

Not applicable.

#### Competing interests

No competing interests.

Received: 25 January 2022 Accepted: 15 June 2022

Published online: 24 June 2022

#### References

- Chen J, Mei K, Xie L, et al. Low vitamin D levels do not aggravate COVID-19 risk or death, and vitamin D supplementation does not improve outcomes in hospitalized patients with COVID-19: a meta-analysis and GRADE assessment of cohort studies and RCTs. *Nutr J*. 2021;20:89. <https://doi.org/10.1186/s12937-021-00744-y>.
- Hastie CE, Mackay DF, Ho F, et al. Vitamin D concentrations and COVID-19 infection in UK Biobank. *Diabetes Metabol Syndr Clin Res Rev*. 2020;14:561–5. <https://doi.org/10.1016/j.dsx.2020.04.050>.
- Hastie CE, Pell JP, Sattar N. Vitamin D and COVID-19 infection and mortality in UK Biobank. *Eur J Nutr*. 2020. <https://doi.org/10.1007/s00394-020-02372-4>.
- Katz J, Yue S, Xue W. Increased risk for Covid-19 in patients with vitamin D deficiency. *Nutrition*. 2020. <https://doi.org/10.1016/j.nut.2020.111106>.
- Kaufman HW, Niles JK, Kroll MH, Bi C, Holick MF. SARS-CoV-2 positivity rates associated with circulating 25-hydroxyvitamin D levels. *PLoS One*. 2020;15(9):e0239252. <https://doi.org/10.1371/journal.pone.0239252>.
- Meltzer DO, Best TJ, Zhang H, Vokes T, Arora V, Solway J. Association of vitamin D status and other clinical characteristics with COVID-19 test results. *JAMA Netw Open*. 2020;3(9):e2019722. <https://doi.org/10.1001/jamanetworkopen.2020.19722>.
- Berry DJ, Dutton J, Fraser WD, Järvelin MR, Hyppönen E. Harmonization study between LC-MS/MS and diasorin RIA for measurement of 25-hydroxyvitamin D concentrations in a large population survey. *J Clin Lab Anal*. 2017;31:e22049. <https://doi.org/10.1002/jcla.22049>.
- Borenstein M, Hedges LV, Higgins JP, Rothstein HR. A basic introduction to fixed-effect and random-effects models for meta-analysis. *Res Synth Methods*. 2010;1(2):97–111. <https://doi.org/10.1002/jrsm.12>.

#### Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

