

LETTER TO THE EDITOR

Open Access



Pharmacological targeting of Tripartite Motif Containing 24 for the treatment of glioblastoma

Mingzhi Han* and Yanfei Sun

Abstract

Glioblastoma (GBM) is the most aggressive brain tumor of the central nervous system. Recent studies have reported the crucial functions of Tripartite Motif Containing 24 (TRIM24) in promoting cancer progression of GBM. However, it remains unclear if TRIM24 is an attractive druggable target for therapeutic intervention in GBM. We therefore performed a series of experiments, aiming to verify whether specific TRIM24 inhibition suppresses GBM malignant functions using dTRIM24 and IACS-9571, two novel selective TRIM24 antagonists. Our data showed that TRIM24 inhibitors serve as effective agents for inhibiting cell propagation and invasion of several patient-derived GBM stem cells (GSCs), and these effects are mediated partially through suppression of the TRIM24-SOX2 axis. This study provides novel insight into the TRIM24-based druggable dependencies, important for developing effective therapeutic strategies for brain tumors.

Keywords: Glioblastoma, TRIM24, Cancer stem cells, Cell viability, Cell Invasion

Letter to the editor

Glioblastoma multiforme (GBM) is the most aggressive brain tumor of the central nervous system. The ability of tumor cells to migrate, rapidly diffuse and invade normal adjacent tissue, their sustained proliferation, and the existence of GBM stem cells (GSCs) leads to a median survival of approximately 15 months following the best standard of care [1–3]. Therefore, it is of paramount importance to understand the molecular mechanisms contributing to GBM development and progression to develop more effective therapies.

Tripartite Motif Containing 24 (TRIM24), also known as TIF1 α , is an important member of the Transcription Intermediary Factor (TIF) family. It consists of a RING-type E3 ubiquitin ligase domain, and a terminal plant homeodomain (PHD)-bromodomain which acts as a reader of the non-canonical histone signature H3K23ac. TRIM24 has been shown to function as an oncogenic

factor or tumor suppressor dependent on the cancer type. For instance, aberrant overexpression of TRIM24 is associated with oncogenesis and disease progression in a wide variety of cancers including breast cancer, gastric cancer, and GBM [4]. Recently, Zhang et al. [5] showed TRIM24 to be highly expressed in GSCs where the binding, through its bromodomain, activates the expression of the pluripotency transcription factor Sex-determining region Y-box 2 (SOX2), -thereby promoting GBM stemness and invasiveness. Through a TRIM24 shRNA knockdown approach and functional assays, it was suggested that TRIM24 represents a potential target for GBM treatment. Recently, potent and specific inhibitors for TRIM24 have been developed [6, 7]. It is therefore important to validate the translational significance of these findings in a pharmacological context.

To determine the antitumor effects of TRIM24-based druggable dependencies, we used two novel TRIM24 inhibitors: (i) IACS-9571, a high-affinity, potent dimethyl-benzimidazole bromodomain inhibitor of TRIM24/BRPF1 with good selectivity over other bromodomain family proteins without modifying TRIM24 expression level [6]; (ii) dTRIM24, a bifunctional

*Correspondence: mingzhi.han@sdu.edu.cn
Cheeloo College of Medicine, Shandong University, Jinan 250012, China



© The Author(s) 2021. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

degrader of TRIM24 based on proteolysis-targeted chimera (PROTAC). dTRIM24 can selectively bind both the bromodomain of TRIM24 and the E3 ubiquitin ligase VHL, thus driving proteasome-mediated degradation of TRIM24 [7]. We determined the effects of these two compounds on a panel of patient-derived GSC lines which have been well characterized (Additional file 1: Table S1). Both dTRIM24 and IACS-9571 effectively and dose-dependently reduced the proliferation of GSCs (Fig. 1A). Furthermore, the treatment of GSCs with dTRIM24 (5, 10 μ M) or IACS-9571 (10, 20 μ M) attenuated the capacity of tumorsphere formation (Fig. 1B and Additional file 1: Fig. S1) and the expression of stemness markers SOX2 and Nestin through immunofluorescence staining (Fig. 1C), demonstrating that pharmacological targeting of TRIM24 effectively inhibits self-renewal of GSCs. Moreover, western blot analysis showed a decrease in TRIM24 and SOX2 expression levels after dTRIM24 treatment (Fig. 1D, upper), verifying its efficacy as a TRIM24 protein degrader in GSCs. Likewise, treatment of IACS-9571 caused a decrease of SOX2 (Fig. 1D, lower). We further observed that both compounds attenuated the invasion distance of GSCs ($P < 0.001$; Fig. 1E) and induced cell apoptosis, while the cell cycle was not significantly affected (Additional file 1: Fig. S2). In a rescue experiment, ectopic expression of SOX2 in GBM#P3 cells partially restored cellular viability suppression followed by IACS-9571 or dTRIM24 treatment compared to the control group (Fig. 1F and Additional file 1: Fig. S3), suggesting that TRIM24-SOX2 axis was involved in the inhibitory effects of these two inhibitors. TRIM24 has been reported to contribute to GBM progression via several signaling pathways. For instance, Zhang et al. [8] found that TRIM24 could bind to the *PIK3CA* (Phosphoinositide-3-Kinase Catalytic Alpha Polypeptide) promoter, thus enhancing PI3K/Akt signaling in GBM cell lines. Lv et al. [4] showed that TRIM24 could cooperatively

activate Signal Transducer and Activator of Transcription 3 (STAT3) signaling and enhance Epidermal Growth Factor Receptor (EGFR)-driven GBM tumorigenesis, indicating multifaceted roles of TRIM24 in the GBM signaling networks. Therefore, further multi-omics studies are warranted in order to elucidate the molecular mechanisms underlying the effects of the TRIM24 inhibitors in GBM.

Conclusions

In conclusion, our data show for the first time that TRIM24 inhibitors serve as effective agents for targeting GSCs, and these inhibitory effects are partially mediated through suppression of the TRIM24-SOX2 axis. These observations, together with the reported ability of dTRIM24 and IACS-9571 to inhibit growth and trigger apoptosis in a panel of acute monocytic leukemia cells [7], make TRIM24 an attractive drug target for therapeutic intervention in GBM. It is also noteworthy that the dTRIM24 is more effective in displacing TRIM24 from chromatin compared to IACS-9571 and exerts a pronounced effect on TRIM24 target genes [7], which is consistent with our findings that dTRIM24 has a relatively lower IC_{50} in GSCs. Based on the Molinspiration Cheminformatics (<http://www.molinspiration.com>) prediction, the Topological Polar Surface Area (TPSA) score of IACS-9571 (TSPA: 122.5) and dTRIM24 (TSPA: 260.0) show that these compounds have relatively moderate to low values of blood-brain barrier (BBB) penetration. This implies that pharmacologically targeting TRIM24 for the treatment of GBM might not, at present, be achieved in a preclinical and clinical context. Yet, optimization of their chemical structures and new therapeutic developments toward TRIM24 warrant further exploitation.

(See figure on next page.)

Fig. 1 **A** IC_{50} curves for dTRIM24 (MedChemExpress, USA) and IACS-9571 (MCE, hydrochloride form) in GBM#P3, GBM#BG7, GBM#06, and GBM#BG5 cells using the Cell Titer-Glo viability assay. **B** Quantification of tumorsphere formation assays for GSCs treated with different concentrations of dTRIM24 (0–10 μ M) (A) or IACS-9571 (0–20 μ M) for 6 days. GSCs (1000 cells/mL/well) were seeded in 6-well ultra-low adhesion plates. Inverted phase-contrast microscopy was used to count the sphere number. **C** Representative images of immunofluorescence staining for SOX2 (red; dilution 1: 100) or Nestin (green; dilution 1: 200) in GBM#P3 treated with dTRIM24 or IACS-9571 for 48 h. Nuclei were counterstained with DAPI (blue). Scale bar = 100 μ m. **D** Western blot analyses of the TRIM24, SOX2, and GAPDH in lysates (20 μ g) from GSCs treated with different concentrations dTRIM24 (0–10 μ M) or IACS-9571 (0–20 μ M). **E** Representative images of spheroids in 3D invasion assays for GBM#P3 GSCs treated with DMSO, dTRIM24 (5 μ M), or IACS-9571 (10 μ M), and evaluated at 24 h. Scale bar = 200 μ m (lower). Graphic representation of the quantification of the distance of invading cells from the tumorspheres determined after 24 h (upper). **F** Relative cell viability for rescue experiments using the Cell Titer-Glo viability assay in GBM#P3 cells as indicated. Data are shown as mean \pm SEM. Statistical significance was determined by one-way ANOVA. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$




Fig. 1 (See legend on previous page.)

Abbreviations

GBM: Glioblastoma; TRIM24: Tripartite Motif Containing 24; TIF: Transcription Intermediary Factor; PHD: Plant homeodomain; GSCs: GBM stem cells; PROTAC: Proteolysis-targeted chimera; SOX2: Sex-determining region Y-box 2; PIK3CA: Phosphoinositide-3-Kinase Catalytic Alpha Polypeptide; STAT3: Signal Transducer and Activator of Transcription 3; EGFR: Epidermal Growth Factor Receptor.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12967-021-03158-w>.

Additional file 1. Additional Figures S1–S3 and Table S1.

Acknowledgements

We thank Dr. Justin Vareecal Joseph for establishing the primary GBM cells.

Authors' contributions

The study design, experiment and performed data analysis, and the writing of the manuscript: MH and YS. All authors read and approved the final manuscript.

Funding

This work was supported by the Shandong Provincial Natural Science Foundation (ZR2021QH030) and the Qilu Young Scholar Program of Shandong University, China.

Availability of data and materials

All data generated or analyzed during this study are included in this article.

Declarations**Ethics approval and consent to participate**

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors have no relevant competing interests to disclose.

Received: 25 August 2021 Accepted: 21 November 2021

Published online: 09 December 2021

References

- Louis DN, Perry A, Reifenberger G, et al. The 2016 World Health Organization Classification of tumors of the central nervous system: a summary. *Acta Neuropathol.* 2016;131(6):803–20.
- Xu X, Wang L, Zang Q, et al. Rewiring of purine metabolism in response to acidosis stress in glioma stem cells. *Cell Death Dis.* 2021;12(3):277.
- Neftel C, Laffy J, Filbin MG, et al. An integrative model of cellular states, plasticity, and genetics for glioblastoma. *Cell.* 2019;178(4):835–849.e21.
- Lv D, Li Y, Zhang W, Alvarez AA, et al. TRIM24 is an oncogenic transcriptional co-activator of STAT3 in glioblastoma. *Nat Commun.* 2017;8(1):1454.
- Zhang LH, Yin YH, Chen HZ, et al. TRIM24 promotes stemness and invasiveness of glioblastoma cells via activating SOX2 expression. *Neuro Oncol.* 2020;22(12):1797–808.
- Palmer WS, Poncet-Montange G, Liu G, et al. Structure-guided design of IACS-9571, a selective high-affinity dual TRIM24-BRPF1 bromodomain inhibitor. *J Med Chem.* 2016;59(4):1440–54.
- Gechijian LN, Buckley DL, Lawlor MA, et al. Functional TRIM24 degrader via conjugation of ineffectual bromodomain and VHL ligands. *Nat Chem Biol.* 2018;14(4):405–12.
- Zhang LH, Yin AA, Cheng JX, et al. TRIM24 promotes glioma progression and enhances chemoresistance through activation of the PI3K/Akt signaling pathway. *Oncogene.* 2015;34(5):600–10.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

